

ARG43889 anti-CD194 / CCR4 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes CD194 / CCR4
Tested Reactivity	Hu, Ms, Rat
Tested Application	ELISA, FACS, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CD194 / CCR4
Species	Human
Immunogen	Human CCR4 recombinant protein
Conjugation	Un-conjugated
Alternate Names	CD194; ChemR13; K5-5; C-C chemokine receptor type 4; CCR-4; HGCN:14099; CKR4; CD antigen CD194; CC-CKR-4; CCR4; C-C CKR-4; CMKBR4

Application Instructions

Application table	Application	Dilution
	ELISA	0.1-0.5 µg/ml
	FACS	1-3 µg/1x10 ⁶ cells
	WB	0.25-0.5 µg/ml

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

Form	Liquid
Purification	Affinity purified with Immunogen.
Buffer	0.9% NaCl, 0.2% Na ₂ HPO ₄ and 4% Trehalose.
Stabilizer	4% Trehalose
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	CCR4
Gene Full Name	C-C Motif Chemokine Receptor 4
Background	The protein encoded by this gene belongs to the G-protein-coupled receptor family . It is a receptor for the CC chemokine - MIP-1, RANTES, TARC and MCP-1. Chemokines are a group of small polypeptide, structurally related molecules that regulate cell trafficking of various types of leukocytes. The chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis.
Function	High affinity receptor for the C-C type chemokines CCL17/TARC, CCL22/MDC and CKLF isoform 1/CKLF1. The activity of this receptor is mediated by G(i) proteins which activate a phosphatidylinositol-calcium second messenger system. Can function as a chemoattractant homing receptor on circulating memory lymphocytes and as a coreceptor for some primary HIV-2 isolates. In the CNS, could mediate hippocampal-neuron survival.
Calculated Mw	41 kDa
PTM	Disulfide bond, Glycoprotein, Phosphoprotein
Cellular Localization	Cell membrane

Images



ARG43889 anti-CD194 / CCR4 antibody WB image

Western blot: 293T and Raji stained with ARG43889 anti-CD194 / CCR4 antibody at 0.5 µg/mL dilution.




ARG43889 anti-CD194 / CCR4 antibody FACS image

Flow Cytometry: HEL cells stained with ARG43889 anti-CD194 / CCR4 antibody (blue) at 1 µg/1x10⁶ cells dilution.

ARG43889 anti-CD194 / CCR4 antibody WB image

Western blot: Rat spleen and Rat thymus with ARG43889 anti-CD194 / CCR4 antibody at 0.5 $\mu\text{g}/\text{mL}$ dilution.



ARG43889 anti-CD194 / CCR4 antibody WB image

Western blot: Mouse spleen and Mouse thymus with ARG43889 anti-CD194 / CCR4 antibody at 0.5 $\mu\text{g}/\text{mL}$ dilution.

