

# Product datasheet

info@arigobio.com

# ARG10242 anti-Insulin antibody [8E2]

Package: 100 μg, 50 μg

Store at: -20°C

#### **Summary**

Product Description Mouse Monoclonal antibody [8E2] recognizes Insulin

Tested Reactivity Hu, Bov, Pig

Tested Application ELISA

Host Mouse

Clonality Monoclonal

Clone 8E2

Isotype IgG1

Target Name Insulin
Species Human

Immunogen purified human insulin

Conjugation Un-conjugated

Alternate Names IDDM; IDDM2; IDDM1; ILPR; MODY10; Insulin; IRDN

## **Application Instructions**

**Application Note** 

\* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

#### **Properties**

Form Liquid

Purification Protein A affinity purified.

Buffer PBS (pH 7.4) and 0.1% Sodium azide

Preservative 0.1% Sodium azide

Concentration 1.0-2.0 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links <u>GeneID: 3630 Human</u>

GeneID: 397415 Pig

#### Swiss-port # P01308 Human

#### Swiss-port # P01315 Pig

Gene Symbol INS

Gene Full Name insulin

Background After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three

peptides: the B chain and A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region.

Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2010]

Function Insulin decreases blood glucose concentration. It increases cell permeability to monosaccharides, amino

acids and fatty acids. It accelerates glycolysis, the pentose phosphate cycle, and glycogen synthesis in

liver. [UniProt]

Research Area Cell Biology and Cellular Response antibody; Metabolism antibody; Neuroscience antibody; Signaling

Transduction antibody

Calculated Mw 12 kDa