

ARG10552 anti-Bax antibody [G26-R]

Package: 50 µl
Store at: -20°C

Summary

Product Description	Rabbit Monoclonal antibody [G26-R] recognizes Bax
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, IP, WB
Host	Rabbit
Clonality	Monoclonal
Clone	G26-R
Isotype	IgG
Target Name	Bax
Species	Human
Immunogen	Synthetic peptide around the N-terminus of Human bax
Epitope	Antibody recognizes the epitope located between Pro13 - Gly36.
Conjugation	Un-conjugated
Alternate Names	Bcl-2-like protein 4; Bcl2-L-4; BCL2L4; Apoptosis regulator BAX

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:100 – 1:600
	IHC-P	1:100
	IP	1:100 – 1:600
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Buffer	20 mM Tris-HCl (pH 8.0), 0.05% Sodium azide and 10 mg/ml BSA
Preservative	0.05% Sodium azide
Stabilizer	10 mg/ml BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 12028 Mouse GeneID: 581 Human Swiss-port # Q07812 Human Swiss-port # Q07813 Mouse
Gene Symbol	BAX
Gene Full Name	BCL2-associated X protein
Background	Bax belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. The association and the ratio of BAX to BCL2 also determines survival or death of a cell following an apoptotic stimulus. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene. [provided by RefSeq, Dec 2019]
Function	Bax plays a role in the mitochondrial apoptotic process. Under normal conditions, BAX is largely cytosolic via constant retrotranslocation from mitochondria to the cytosol mediated by BCL2L1/Bcl-xL, which avoids accumulation of toxic BAX levels at the mitochondrial outer membrane (MOM) (PubMed:21458670). Under stress conditions, undergoes a conformation change that causes translocation to the mitochondrion membrane, leading to the release of cytochrome c that then triggers apoptosis. Promotes activation of CASP3, and thereby apoptosis. [UniProt]
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Cell Death antibody; Metabolism antibody; Mitochondrial fission antibody; Apoptosis Marker antibody; Pro-apoptotic Bcl2 protein antibody
Calculated Mw	21 kDa

Images



ARG10552 anti-Bax antibody [G26-R] WB image

Western blot: 20 μ l of 293T, Mouse brain and Rat brain lysates stained with ARG10552 anti-Bax antibody [G26-R] at 1:1000 dilution.



ARG10552 anti-Bax antibody [G26-R] ICC/IF image

Immunocytochemistry: HEK293 cells stained with ARG10552 anti-Bax antibody [G26-R] at 1:100 dilution.