

Product datasheet

info@arigobio.com

ARG10847 anti-Complement C3 antibody [E28-P]

Package: 100 μl Store at: -20°C

Summary

Host

Product Description Rabbit Monoclonal antibody [E28-P] recognizes Complement C3

Tested Reactivity Hu
Tested Application IHC-P

Clonality Monoclonal

Clone E28-P

Target Name Complement C3

Rabbit

Species Human

Immunogen Synthetic peptide around the N-terminus of Human Complement C3.

Conjugation Un-conjugated

Alternate Names CPAMD1; HEL-S-62p; ASP; ARMD9; C3a; C3b; C3bc; Complement C3; C3adesArg; AHUS5; C3 and PZP-

like alpha-2-macroglobulin domain-containing protein 1

Application Instructions

Application table	Application	Dilution
	IHC-P	1:100 - 1:200
Application Note	IHC-P: Antigen Retrieval: Boil tissue section in Tris-EDTA buffer (pH 9.0) for 40 min followed by cooling at RT for 20 min.	
	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	
	should be determined by the scientist.	

Properties

Form Liquid

Buffer 20 mM Tris-HCl (pH 8.0), 0.05% Sodium azide and 20 mg/ml BSA.

Preservative 0.05% Sodium azide

Stabilizer 20 mg/ml BSA

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol C3

Gene Full Name complement component 3

Background Complement component C3 plays a central role in the activation of complement system. Its activation

is required for both classical and alternative complement activation pathways. A peptide (C3a) derived from the encoded protein has antimicrobial activity, so people with C3 deficiency are susceptible to

bacterial infection. [provided by RefSeq, Nov 2014]

Function C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can

bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

Derived from proteolytic degradation of complement C3, C3a anaphylatoxin is a mediator of local inflammatory process. In chronic inflammation, acts as a chemoattractant for neutrophils (By similarity). It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes.

C3-beta-c: Acts as a chemoattractant for neutrophils in chronic inflammation.

Acylation stimulating protein: adipogenic hormone that stimulates triglyceride (TG) synthesis and glucose transport in adipocytes, regulating fat storage and playing a role in postprandial TG clearance. Appears to stimulate TG synthesis via activation of the PLC, MAPK and AKT signaling pathways. Ligand for C5AR2. Promotes the phosphorylation, ARRB2-mediated internalization and recycling of C5AR2.

[UniProt]

Calculated Mw 187 kDa


PTM C3b is rapidly split in two positions by factor I and a cofactor to form iC3b (inactivated C3b) and C3f

which is released. Then iC3b is slowly cleaved (possibly by factor I) to form C3c (beta chain + alpha' chain fragment 1 + alpha' chain fragment 2), C3dg and C3f. Other proteases produce other fragments such as C3d or C3g. C3a is further processed by carboxypeptidases to release the C-terminal arginine residue generating the acylation stimulating protein (ASP). Levels of ASP are increased in adipocytes in

the postprandial period and by insulin and dietary chylomicrons.

Phosphorylated by FAM20C in the extracellular medium. [UniProt]

Images



ARG10847 anti-Complement C3 antibody [E28-P] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human skin biopsy from the lesion of the early pemphigus vulgaris (without blister formation) (4 μ m section) stained with ARG10847 anti-Complement C3 antibody [E28-P].