

ARG20922 anti-CD38 antibody [NIMR-5] (PE-Cyanine 5)

Package: 50 µg
Store at: 4°C

Summary

Product Description	PE-Cyanine 5-conjugated Rat Monoclonal antibody [NIMR-5] recognizes CD38
Tested Reactivity	Ms
Tested Application	BL, Cell-Act , FACS, IHC-Fr
Specificity	Mouse CD38.
Host	Rat
Clonality	Monoclonal
Clone	NIMR-5
Isotype	IgG2a, kappa
Target Name	CD38
Species	Mouse
Immunogen	BCL1 plasma membrane glycoproteins
Conjugation	PE-Cyanine 5
Alternate Names	cADPr hydrolase 1; ADPRC 1; EC 3.2.2.6; 2'-phospho-ADP-ribose cyclase/2'-phospho-cyclic-ADP-ribose transferase; Cyclic ADP-ribose hydrolase 1; ADPRC1; EC 2.4.99.20; ADP-ribose cyclase 1; 2'-phospho-cyclic-ADP-ribose transferase; CD antigen CD38; T10; 2'-phospho-ADP-ribose cyclase; ADP-ribose cyclase/cyclic ADP-ribose hydrolase 1

Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	Cell-Act	Assay-dependent
	FACS	< 0.2 µg/10 ⁶ cells
	IHC-Fr	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties


Form	Liquid
Buffer	PBS, 0.1% Sodium azide and Sucrose.
Preservative	0.1% Sodium azide
Stabilizer	Sucrose
Concentration	0.1 mg/ml

Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 12494 Mouse Swiss-port # P56528 Mouse
Gene Symbol	CD38
Gene Full Name	CD38 antigen
Background	The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]
Function	Synthesizes the second messengers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system. [UniProt]
Research Area	Cancer antibody; Developmental Biology antibody; Immune System antibody; Metabolism antibody; Pro-B Cell Marker antibody; Pre-B Cell Marker antibody
Calculated Mw	34 kDa

Images



ARG20922 anti-CD38 antibody [NIMR-5] (PE-Cyanine 5) FACS image

Flow Cytometry: BALB/c Mouse splenocytes stained with ARG20922 anti-CD38 antibody [NIMR-5] (PE-Cyanine 5) and [ARG65534](#) anti-CD45R / B220 antibody [RA3-6B2] (PE).