

## ARG20947 anti-CD72 antibody [10.1.D2]

Package: 250 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [10.1.D2] recognizes CD72
Tested Reactivity	Ms
Tested Application	BL, Cell-Act , FACS, IP
Specificity	Mouse CD72.1. The clone 10.1.D2 blocks binding of CD5 to CD72 on the B cell surface which leads to inhibition of the positive signal resulting from CD5/CD72 pairing. However, 10.1.D2 can itself trigger a positive signal by binding CD72.
Host	Mouse
Clonality	Monoclonal
Clone	10.1.D2
Isotype	IgG2b, kappa
Target Name	CD72
Species	Mouse
Immunogen	DBA/2 mouse spleen cells
Conjugation	Un-conjugated
Alternate Names	CD antigen CD72; LYB2; B-cell differentiation antigen CD72; Lyb-2; CD72b

### Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	Cell-Act	Assay-dependent
	FACS	Assay-dependent
	IP	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

**Note**

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links	<a href="#">GeneID: 12517 Mouse</a> <a href="#">Swiss-port # P21855 Mouse</a>
Gene Symbol	CD72
Gene Full Name	CD72 antigen
Function	Plays a role in B-cell proliferation and differentiation. [UniProt]
Calculated Mw	40 kDa
PTM	Phosphorylated upon engagement of the B-cell receptor, probably by LYN or SYK. Phosphorylation at Tyr-7 is important for interaction with PTPN6/SHP-1 (By similarity).