

## ARG21039 anti-CD81 antibody [2F7] (PE)

Package: 50 µg  
Store at: 4°C

### Summary

Product Description	PE-conjugated Hamster Monoclonal antibody [2F7] recognizes CD81
Tested Reactivity	Ms
Tested Application	BL, FACS, ICC/IF, IHC-P
Specificity	Mouse CD81. The clone 2F7 can block thymocyte interaction with CD81 in vitro.
Host	Hamster
Clonality	Monoclonal
Clone	2F7
Isotype	IgG3
Target Name	CD81
Species	Mouse
Immunogen	Mouse epithelial cell line PAM212
Conjugation	PE
Alternate Names	CD antigen CD81; TAPA1; Tspan-28; S5.7; CD81 antigen; Target of the antiproliferative antibody 1; Tetraspanin-28; 26 kDa cell surface protein TAPA-1; CVID6; TSPAN28

### Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	FACS	< 0.2 µg/10 <sup>6</sup> cells
	ICC/IF	Assay-dependent
	IHC-P	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Buffer	PBS, 0.1% Sodium azide and Sucrose.
Preservative	0.1% Sodium azide
Stabilizer	Sucrose
Concentration	0.1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note


For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links	<a href="#">GeneID: 12520 Mouse</a> <a href="#">Swiss-port # P35762 Mouse</a>
Gene Symbol	CD81
Gene Full Name	CD81 antigen
Background	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins. This protein appears to promote muscle cell fusion and support myotube maintenance. Also it may be involved in signal transduction. This gene is localized in the tumor-suppressor gene region and thus it is a candidate gene for malignancies. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2014]
Function	May play an important role in the regulation of lymphoma cell growth. Interacts with a 16-kDa Leu-13 protein to form a complex possibly involved in signal transduction. May act as the viral receptor for HCV. [UniProt]
Highlight	Related products: <a href="#">CD81 antibodies: Anti-Hamster IgG secondary antibodies:</a> Related news: <a href="#">Tools for studying Exosomes</a>
Calculated Mw	26 kDa
PTM	Not glycosylated.

## Images



ARG21039 anti-CD81 antibody [2F7] (PE) FACS image

Flow Cytometry: Hamster CHO-K1 cell line and BALB/cAnN Mouse B lymphocyte cell line A20 stained with ARG21039 anti-CD81 antibody [2F7] (PE).