

ARG21182 anti-CD28 antibody [AV7]

Package: 100 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [AV7] recognizes CD28
Tested Reactivity	Chk, Turkey
Tested Application	FACS, ICC/IF, IHC-Fr, IP
Specificity	Chicken/Turkey CD28. The clone AV7 also reacts with turkey CD28 as demonstrated by flow cytometry.
Host	Mouse
Clonality	Monoclonal
Clone	AV7
Isotype	IgG1, kappa
Target Name	CD28
Species	Chicken
Immunogen	Chicken peripheral blood lymphocytes
Conjugation	Un-conjugated
Alternate Names	Tp44; TP44; T-cell-specific surface glycoprotein CD28; CD antigen CD28

Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
	IP	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 396249 Chicken Swiss-port # P31043 Chicken
Gene Symbol	CD28
Gene Full Name	CD28 molecule
Background	The protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2 development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]
Function	Involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival. [UniProt]
Research Area	Developmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibody
Calculated Mw	25 kDa
PTM	CD40LG induces tyrosine phosphorylation of isoform 3.