

Product datasheet

info@arigobio.com

ARG21233 anti-CD44 antibody [AV6] (FITC)

Package: 100 μg Store at: 4°C

Summary

Product Description FITC-conjugated Mouse Monoclonal antibody [AV6] recognizes CD44

Tested Reactivity Chk, Turkey

Tested Application FACS, IHC-Fr

Specificity Chicken/Turkey CD44. The clone AV6 also reacts with turkey peripheral blood mononuclear cells.

Host Mouse

Clonality Monoclonal

Clone AV6

Isotype IgG1, kappa

Target Name CD44
Species Chicken
Conjugation FITC

Alternate Names MDU2; MDU3; GP90 lymphocyte homing/adhesion receptor; Hermes antigen; Extracellular matrix

receptor III; PGP-I; Epican; CDW44; Phagocytic glycoprotein 1; Pgp1; HUTCH-I; MC56; Hyaluronate receptor; CD antigen CD44; Heparan sulfate proteoglycan; CD44 antigen; LHR; IN; HCELL; Phagocytic

glycoprotein I; PGP-1; CSPG8; MIC4; ECMR-III; CDw44

Application Instructions

Application table	Application	Dilution
	FACS	< 1 μg/10^6 cells
	IHC-Fr	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Buffer PBS and 0.1% Sodium azide.

Preservative 0.1% Sodium azide

Concentration 0.5 mg/ml

Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol

CD44

Gene Full Name

CD44 molecule (Indian blood group)

Background

The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis. [provided by RefSeq, Jul 2008]

Function

Receptor for hyaluronic acid (HA). Mediates cell-cell and cell-matrix interactions through its affinity for HA, and possibly also through its affinity for other ligands such as osteopontin, collagens, and matrix metalloproteinases (MMPs). Adhesion with HA plays an important role in cell migration, tumor growth and progression. In cancer cells, may play an important role in invadopodia formation. Also involved in lymphocyte activation, recirculation and homing, and in hematopoiesis. Altered expression or dysfunction causes numerous pathogenic phenotypes. Great protein heterogeneity due to numerous alternative splicing and post-translational modification events. [UniProt]

Research Area

Cancer antibody; Developmental Biology antibody; Immune System antibody; Chondrogenesis Study antibody

Calculated Mw

82 kDa


PTM

Proteolytically cleaved in the extracellular matrix by specific proteinases (possibly MMPs) in several cell lines and tumors.

N- and O-glycosylated. O-glycosylation contains more-or-less-sulfated chondroitin sulfate glycans, whose number may affect the accessibility of specific proteinases to their cleavage site(s). It is uncertain if O-glycosylation occurs on Thr-637 or Thr-638.


Phosphorylated; activation of PKC results in the dephosphorylation of Ser-706 (constitutive phosphorylation site), and the phosphorylation of Ser-672.

Images



ARG21233 anti-CD44 antibody [AV6] (FITC) FACS image

Flow Cytometry: Chicken peripheral blood mononuclear cells stained with ARG21233 anti-CD44 antibody [AV6] (FITC).



ARG21233 anti-CD44 antibody [AV6] (FITC) FACS image

Flow Cytometry: Chicken peripheral blood mononuclear cells stained with $\underline{\mathsf{ARG21246}}$ anti-Monocyte + Macrophage antibody [KUL01] (PE) and $\underline{\mathsf{ARG21233}}$ anti-CD44 antibody [AV6] (FITC).