

Product datasheet

info@arigobio.com

ARG21333 anti-CD14 antibody [UCHM-1] (PE-Cyanine 5)

Package: 50 tests Store at: 4°C

Summary

Product Description PE-Cyanine 5-conjugated Mouse Monoclonal antibody [UCHM-1] recognizes CD14

Tested Reactivity Hu

Tested Application BL, Depletion, FACS, ICC/IF, IHC-Fr

Specificity Human/Cynomolgus CD14.

Host Mouse

Clonality Monoclonal

Clone UCHM-1

Isotype IgG2a, kappa

Target Name CD14

Species Human

Immunogen Thymocytes and peripheral blood lymphocytes from a remission AML patient

Conjugation PE-Cyanine 5

Alternate Names CD antigen CD14; Myeloid cell-specific leucine-rich glycoprotein; Monocyte differentiation antigen

CD14

Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	Depletion	Assay-dependent
	FACS	10 μl/10^6 cells
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid	
Buffer	PBS, 0.1% Sodium azide and Sucrose.	
Preservative	0.1% Sodium azide	
Stabilizer	Sucrose	
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be	

gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links <u>GeneID: 929 Human</u>

Swiss-port # P08571 Human

Gene Symbol CD14

Gene Full Name CD14 molecule

Background The protein encoded by this gene is a surface antigen that is preferentially expressed on

monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide. Alternative splicing results in multiple transcript variants encoding the

same protein. [provided by RefSeq, Mar 2010]

Function In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the MD-2/TLR4 complex,

thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response.

Up-regulates cell surface molecules, including adhesion molecules. [UniProt]


Research Area Developmental Biology antibody; Immune System antibody; General Lymphocyte Marker Study

antibody; Macrophages and neutrophils antibody

Calculated Mw 40 kDa

PTM N- and O- glycosylated. O-glycosylated with a core 1 or possibly core 8 glycan.

Images



ARG21333 anti-CD14 antibody [UCHM-1] (PE-Cyanine 5) FACS image

Flow Cytometry: Human peripheral blood monocytes and lymphocytes stained with ARG21333 anti-CD14 antibody [UCHM-1] (PE-Cyanine 5) and Mouse anti-Human CD3 antibody (PE).