

ARG21438 anti-IL4 antibody [MP4-25D2] (low endotoxin)

Package: 100 µg
Store at: -20°C

Summary

Product Description	Azide free and low endotoxin Rat Monoclonal antibody [MP4-25D2] recognizes IL4
Tested Reactivity	Hu, Bb, R. Mk
Tested Application	ELISA, ELISPOT, FACS, IHC-Fr, Neut
Specificity	Human/Rhesus/Cynomolgus/Baboon IL-4.
Host	Rat
Clonality	Monoclonal
Clone	MP4-25D2
Isotype	IgG1, kappa
Target Name	IL4
Species	Human
Immunogen	CHO-expressed human IL-4
Conjugation	Un-conjugated
Alternate Names	BSF-1; IL-4; B-cell stimulatory factor 1; BCGF1; Lymphocyte stimulatory factor 1; BCGF-1; Interleukin-4; Pitrakinra; Binetrakin; BSF1

Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	ELISPOT	Assay-dependent
	FACS	Assay-dependent
	IHC-Fr	Assay-dependent
	Neut	Assay-dependent

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification Note	Low endotoxin
Buffer	PBS
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw

cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links

[GeneID: 3565 Human](#)

[Swiss-port # P05112 Human](#)

Gene Symbol

IL4

Gene Full Name

interleukin 4

Background

The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Function

Participates in at least several B-cell activation processes as well as of other cell types. It is a costimulator of DNA-synthesis. It induces the expression of class II MHC molecules on resting B-cells. It enhances both secretion and cell surface expression of IgE and IgG1. It also regulates the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. [UniProt]

Calculated Mw

17 kDa