

## Product datasheet

info@arigobio.com

# ARG21488 anti-IL17 / IL17A antibody [TC11-8H4] (Biotin)

Package: 100 μg Store at: 4°C

#### **Summary**

Product Description Biotin-conjugated Rat Monoclonal antibody [TC11-8H4 ] recognizes IL17 / IL17A

Tested Reactivity Ms

Tested Application ELISA, ELISPOT, FACS

Specificity Mouse IL-17A.

Host Rat

**Clonality** Monoclonal

Clone TC11-8H4

Isotype IgG1, kappa
Target Name IL17 / IL17A

Species Mouse

Immunogen E. coli-expressed mouse IL-17

Conjugation Biotin

Alternate Names IL-17A; Interleukin-17A; IL-17; CTLA8; IL17; CTLA-8; Cytotoxic T-lymphocyte-associated antigen 8

#### **Application Instructions**

Application table	Application	Dilution
	ELISA	1:1000 - 1:2000
	ELISPOT	Assay-dependent
	FACS	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### **Properties**

Form Liquid

Buffer PBS and 0.1% Sodium azide.

Preservative 0.1% Sodium azide

Concentration 0.5 mg/ml

Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Database links <u>GeneID: 16171 Mouse</u>

Swiss-port # Q62386 Mouse

Gene Symbol IL17A

Gene Full Name interleukin 17A

Background The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This

cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can

stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. [provided by

RefSeq, Jul 2008]

Function Induces stromal cells to produce proinflammatory and hematopoietic cytokines. Enhances the surface

expression of ICAM1/intracellular adhesion molecule 1 in fibroblasts. [UniProt]

Highlight Related products:

IL17 antibodies; IL17 ELISA Kits; IL17 recombinant proteins; Anti-Rat IgG secondary antibodies;

Related news:

HMGB1, a biomarker and therapeutic target in COVID-19

Calculated Mw 18 kDa

PTM Found both in glycosylated and nonglycosylated forms.