

ARG21530 Goat anti-Mouse IgM antibody (TRITC), pre-adsorbed

Package: 500 µg
Store at: 4°C

Summary

| | |
|---------------------|--|
| Product Description | TRITC-conjugated Goat Polyclonal antibody recognizes Mouse IgM |
| Tested Reactivity | Ms |
| Tested Application | FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB |
| Specificity | The antibody reacts with the heavy chain of Mouse IgM. The antibody is pre-adsorbed with Mouse IgG and IgA, so the antibody may not react with Mouse IgG and IgA, but may react with immunoglobulins from other species. |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | IgM |
| Species | Mouse |
| Conjugation | TRITC |

Application Instructions

| Pre Adsorbed | Mouse IgG and IgA. | | | | | | | | | | | | | | |
|-------------------|---|-------------|----------|------|-----------------|-------|---------------|--------|-----------------|--------|-----------------|-------|-----------------|----|-----------------|
| Application table | <table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>1:100 - 1:400</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table> | Application | Dilution | FACS | Assay-dependent | FLISA | 1:100 - 1:400 | ICC/IF | Assay-dependent | IHC-Fr | Assay-dependent | IHC-P | Assay-dependent | WB | Assay-dependent |
| Application | Dilution | | | | | | | | | | | | | | |
| FACS | Assay-dependent | | | | | | | | | | | | | | |
| FLISA | 1:100 - 1:400 | | | | | | | | | | | | | | |
| ICC/IF | Assay-dependent | | | | | | | | | | | | | | |
| IHC-Fr | Assay-dependent | | | | | | | | | | | | | | |
| IHC-P | Assay-dependent | | | | | | | | | | | | | | |
| WB | Assay-dependent | | | | | | | | | | | | | | |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | | | | | | | | | | | | | | |

Properties

| | |
|---------------|---------------------------------------|
| Form | Liquid |
| Purification | Affinity purification with immunogen. |
| Buffer | PBS and 0.1% Sodium azide. |
| Preservative | 0.1% Sodium azide |
| Concentration | 1 mg/ml |


Storage instruction

Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Images



ARG21530 Goat anti-Mouse IgM antibody (TRITC) (pre-adsorbed)
FLISA image

FLISA: The plate was coated with purified Mouse IgM, IgG and IgA. Immunoglobulins were detected with serially diluted ARG21530 Goat anti-Mouse IgM antibody (TRITC) (pre-adsorbed).