

ARG21557

Goat anti-Mouse Lambda Light Chain antibody (TRITC), pre-adsorbed

Package: 500 µg

Store at: 4°C

Summary

Product Description	TRITC-conjugated Goat Polyclonal antibody recognizes Mouse Lambda Light Chain
Tested Reactivity	Ms
Tested Application	FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB
Specificity	The antibody reacts with Mouse lambda light chains. The antibody is pre-adsorbed with Mouse kappa light chains, so the antibody may not react with Mouse kappa light chains, but may react with lambda light chains from other species.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	Lambda Light Chain
Species	Mouse
Conjugation	TRITC

Application Instructions

Pre Adsorbed	Mouse kappa light chains.														
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>1:100 - 1:400</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	FACS	Assay-dependent	FLISA	1:100 - 1:400	ICC/IF	Assay-dependent	IHC-Fr	Assay-dependent	IHC-P	Assay-dependent	WB	Assay-dependent
Application	Dilution														
FACS	Assay-dependent														
FLISA	1:100 - 1:400														
ICC/IF	Assay-dependent														
IHC-Fr	Assay-dependent														
IHC-P	Assay-dependent														
WB	Assay-dependent														
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.														

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	1 mg/ml


Storage instruction

Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Images



ARG21557 Goat anti-Mouse Lambda Light Chain antibody (TRITC) (pre-adsorbed) IHC-Fr image

Immunohistochemistry: Frozen section of Mouse lymph node tissue stained with ARG21557 Goat anti-Mouse Lambda Light Chain antibody (TRITC) (pre-adsorbed) followed by DAPI.