

ARG21581 Goat anti-Mouse IgE antibody, pre-adsorbed

Package: 500 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes Mouse IgE
Tested Reactivity	Ms
Tested Application	ELISA, FACS, FLISA, IHC-P, IP, WB
Specificity	The antibody reacts with the heavy chain of Mouse IgE. The antibody is pre-adsorbed with Mouse IgG1, IgG2a, IgG2b, IgG3, IgM and IgA, so the antibody may not react with Mouse IgG1, IgG2a, IgG2b, IgG3, IgM and IgA, but may react with immunoglobulins from other species.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgE
Species	Mouse
Conjugation	Un-conjugated

Application Instructions

Pre Adsorbed	Mouse IgG1, IgG2a, IgG2b, IgG3, IgM and IgA.														
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>IP</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	ELISA	Assay-dependent	FACS	Assay-dependent	FLISA	Assay-dependent	IHC-P	Assay-dependent	IP	Assay-dependent	WB	Assay-dependent
Application	Dilution														
ELISA	Assay-dependent														
FACS	Assay-dependent														
FLISA	Assay-dependent														
IHC-P	Assay-dependent														
IP	Assay-dependent														
WB	Assay-dependent														
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.														

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	BBS (pH 8.2)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.