

ARG21652 Goat anti-Human IgM antibody, F(ab')₂ fragment, pre-adsorbed

Package: 500 µg
Store at: -20°C

Summary

Product Description	F(ab') ₂ fragment of Goat Polyclonal antibody recognizes Human IgM
Tested Reactivity	Hu
Tested Application	ELISA, ELISPOT, FACS, FLISA, WB
Specificity	The antibody reacts with the heavy chain of Human IgM. The antibody is pre-adsorbed with Human IgG and IgA, so the antibody may not react with Human IgG and IgA, but may react with IgM from other species.
Host	Goat
Clonality	Polyclonal
Isotype	F(ab') ₂ IgG
Target Name	IgM
Species	Human
Conjugation	Un-conjugated

Application Instructions

Pre Adsorbed	Human IgG and IgA.												
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>Assay-dependent</td></tr><tr><td>ELISPOT</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	ELISA	Assay-dependent	ELISPOT	Assay-dependent	FACS	Assay-dependent	FLISA	Assay-dependent	WB	Assay-dependent
Application	Dilution												
ELISA	Assay-dependent												
ELISPOT	Assay-dependent												
FACS	Assay-dependent												
FLISA	Assay-dependent												
WB	Assay-dependent												
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.												

Properties


Form	Liquid
Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Highlight

Related news:

[Comprehensive anti-Human secondary antibodies](#)

Images



ARG21652 Goat anti-Human IgM antibody, F(ab')₂ fragment (pre-adsorbed) ELISA image

ELISA: The plate was coated with purified Human IgM, IgG, and IgA. Immunoglobulins were detected with serially diluted ARG21652 Goat anti-Human IgM antibody, F(ab')₂ fragment (pre-adsorbed) followed by [ARG23857](#) Swine anti-Goat IgG (H+L) antibody (HRP) (pre-adsorbed).