

ARG21656

Goat anti-Human IgM antibody, F(ab')₂ fragment (low endotoxin), pre-adsorbed

Package: 250 µg

Store at: -20°C

Summary

Product Description	Azide free and low endotoxin F(ab') ₂ fragment of Goat Polyclonal antibody recognizes Human IgM
Tested Reactivity	Hu
Tested Application	ELISA, ELISPOT, FACS, FLISA
Specificity	The antibody reacts with the heavy chain of Human IgM. The antibody is pre-adsorbed with Human IgG and IgA, so the antibody may not react with Human IgG and IgA, but may react with IgM from other species.
Host	Goat
Clonality	Polyclonal
Isotype	F(ab') ₂ IgG
Target Name	IgM
Species	Human
Conjugation	Un-conjugated

Application Instructions

Pre Adsorbed	Human IgG and IgA.										
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>Assay-dependent</td></tr><tr><td>ELISPOT</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	ELISA	Assay-dependent	ELISPOT	Assay-dependent	FACS	Assay-dependent	FLISA	Assay-dependent
Application	Dilution										
ELISA	Assay-dependent										
ELISPOT	Assay-dependent										
FACS	Assay-dependent										
FLISA	Assay-dependent										
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.										

Properties

Form	Liquid
Purification Note	Low endotoxin
Buffer	PBS
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.