

ARG21676 Goat anti-Human IgA antibody (PE), pre-adsorbed

Package: 250 µg
Store at: 4°C

Summary

| | |
|---------------------|--|
| Product Description | PE-conjugated Goat Polyclonal antibody recognizes Human IgA |
| Tested Reactivity | Hu |
| Tested Application | Depletion, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB |
| Specificity | The antibody reacts with the heavy chain of Human IgA. The antibody is pre-adsorbed with Human IgG and IgM, so the antibody may not react with Human IgG and IgM, but may react with IgA from other species. |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | IgA |
| Species | Human |
| Conjugation | PE |

Application Instructions

Pre Adsorbed Human IgG and IgM.

Application table

| Application | Dilution |
|-------------|--------------------------------|
| Depletion | Assay-dependent |
| FACS | < 0.1 µg/10 ⁶ cells |
| FLISA | < 1 µg/ml |
| ICC/IF | Assay-dependent |
| IHC-Fr | Assay-dependent |
| IHC-P | Assay-dependent |
| WB | Assay-dependent |

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties


| | |
|--------------|---------------------------------------|
| Form | Liquid |
| Purification | Affinity purification with immunogen. |
| Buffer | PBS, 0.1% Sodium azide and Sucrose. |
| Preservative | 0.1% Sodium azide |

| | |
|---------------------|--|
| Stabilizer | Sucrose |
| Concentration | 0.5 mg/ml |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |

Bioinformation

Highlight Related news:
[Comprehensive anti-Human secondary antibodies](#)

Images



ARG21676 Goat anti-Human IgA antibody (PE) (pre-adsorbed) FLISA image

FLISA: The plate was coated with purified Human IgA, IgG and IgM. Immunoglobulins were detected with serially diluted ARG21676 Goat anti-Human IgA antibody (PE) (pre-adsorbed).