

ARG21684 Goat anti-Rat IgM antibody (AP), pre-adsorbed

Package: 500 µl
Store at: 4°C

Summary

Product Description	AP-conjugated Goat Polyclonal antibody recognizes Rat IgM
Tested Reactivity	Rat
Tested Application	ELISA, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB
Specificity	The antibody reacts with the heavy chain of Rat IgM. The antibody is pre-adsorbed with Rat IgG, so the antibody may not react with Rat IgG, but may react with immunoglobulins from other species.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgM
Species	Rat
Conjugation	AP

Application Instructions

Pre Adsorbed	Rat IgG.																
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>1:2000 - 1:4000</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	ELISA	1:2000 - 1:4000	FACS	Assay-dependent	FLISA	Assay-dependent	ICC/IF	Assay-dependent	IHC-Fr	Assay-dependent	IHC-P	Assay-dependent	WB	Assay-dependent
Application	Dilution																
ELISA	1:2000 - 1:4000																
FACS	Assay-dependent																
FLISA	Assay-dependent																
ICC/IF	Assay-dependent																
IHC-Fr	Assay-dependent																
IHC-P	Assay-dependent																
WB	Assay-dependent																
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.																

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	50 mM Tris (pH 8.0), 0.1% Sodium azide, 1 mM MgCl ₂ and 50% Glycerol
Preservative	0.1% Sodium azide
Stabilizer	50% Glycerol


Storage instruction

Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Images



ARG21684 Goat anti-Rat IgM antibody (AP) (pre-adsorbed) ELISA image

ELISA: The plate was coated with purified Rat IgM and IgG. Immunoglobulins were detected with serially diluted ARG21684 Goat anti-Rat IgM antibody (AP) (pre-adsorbed).