

ARG21724
Goat anti-Rabbit IgG antibody (AP), pre-adsorbedPackage: 500 µl
Store at: 4°C

Summary

Product Description	AP-conjugated Goat Polyclonal antibody recognizes Rabbit IgG
Tested Reactivity	Rb
Tested Application	ELISA, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB
Specificity	The antibody reacts with the heavy chain of Rabbit IgG. The antibody is pre-adsorbed with Rabbit IgM, so the antibody may not react with Rabbit IgM, but may react with immunoglobulins from other species.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgG
Species	Rabbit
Conjugation	AP

Application Instructions


Pre Adsorbed	Rabbit IgM.																
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>1:2000 - 1:4000</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	ELISA	1:2000 - 1:4000	FACS	Assay-dependent	FLISA	Assay-dependent	ICC/IF	Assay-dependent	IHC-Fr	Assay-dependent	IHC-P	Assay-dependent	WB	Assay-dependent
Application	Dilution																
ELISA	1:2000 - 1:4000																
FACS	Assay-dependent																
FLISA	Assay-dependent																
ICC/IF	Assay-dependent																
IHC-Fr	Assay-dependent																
IHC-P	Assay-dependent																
WB	Assay-dependent																
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.																

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	50 mM Tris (pH 8.0), 0.1% Sodium azide, 1 mM MgCl ₂ and 50% Glycerol
Preservative	0.1% Sodium azide

Stabilizer	50% Glycerol
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Images



ARG21724 Goat anti-Rabbit IgG antibody (AP) (pre-adsorbed) ELISA image

ELISA: The plate was coated with purified rabbit IgG and IgM. Immunoglobulins were detected with serially diluted ARG21724 Goat anti-Rabbit IgG antibody (AP) (pre-adsorbed).