

ARG21913 Goat anti-Human IgA antibody, F(ab')₂ fragment, pre-adsorbed

Package: 250 µg
Store at: -20°C

Summary

Product Description	F(ab') ₂ fragment of Goat Polyclonal antibody recognizes Human IgA
Tested Reactivity	Hu
Tested Application	ELISA, ELISPOT, FACS, FLISA, IHC-Fr
Specificity	The antibody reacts with the heavy chain of Human IgA. The antibody is pre-adsorbed with Human IgG and IgM, so the antibody may not react with Human IgG and IgM, but may react with IgA from other species.
Host	Goat
Clonality	Polyclonal
Isotype	F(ab') ₂ IgG
Target Name	IgA
Species	Human
Conjugation	Un-conjugated

Application Instructions

Pre Adsorbed	Human IgG and IgM.												
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>Assay-dependent</td></tr><tr><td>ELISPOT</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	ELISA	Assay-dependent	ELISPOT	Assay-dependent	FACS	Assay-dependent	FLISA	Assay-dependent	IHC-Fr	Assay-dependent
Application	Dilution												
ELISA	Assay-dependent												
ELISPOT	Assay-dependent												
FACS	Assay-dependent												
FLISA	Assay-dependent												
IHC-Fr	Assay-dependent												
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.												

Properties

Form	Liquid
Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Highlight

Related news:

[Comprehensive anti-Human secondary antibodies](#)