

## ARG21914

### Goat anti-Human IgA antibody, F(ab')<sub>2</sub> fragment (FITC), pre-adsorbed

Package: 250 µg

Store at: 4°C

#### Summary

|                     |  |
|---------------------|--|
| Product Description | FITC-conjugated F(ab') <sub>2</sub> fragment of Goat Polyclonal antibody recognizes Human IgA  |
| Tested Reactivity   | Hu   |
| Tested Application  | FACS, FLISA, IHC-Fr  |
| Specificity         | The antibody reacts with the heavy chain of Human IgA. The antibody is pre-adsorbed with Human IgG and IgM, so the antibody may not react with Human IgG and IgM, but may react with IgA from other species. |
| Host                | Goat   |
| Clonality           | Polyclonal   |
| Isotype             | F(ab') <sub>2</sub> IgG  |
| Target Name         | IgA  |
| Species             | Human  |
| Conjugation         | FITC   |

#### Application Instructions

| Pre Adsorbed      | Human IgG and IgM.   |                              |
|-------------------|--|------------------------------|
| Application table | Application  | Dilution                     |
|                   | FACS   | < 1 µg/10 <sup>6</sup> cells |
|                   | FLISA  | 1:100 - 1:400                |
|                   | IHC-Fr   | Assay-dependent              |
| Application Note  | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. |                              |


#### Properties

|                     |  |
|---------------------|--|
| Form                | Liquid   |
| Buffer              | PBS and 0.1% Sodium azide.   |
| Preservative        | 0.1% Sodium azide  |
| Concentration       | 0.5 mg/ml  |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note                | For laboratory research only, not for drug, diagnostic or other use.   |

Highlight

Related news:  
[Comprehensive anti-Human secondary antibodies](#)

Images



ARG21914 Goat anti-Human IgA antibody, F(ab')<sub>2</sub> fragment (FITC) (pre-adsorbed) FLISA image

FLISA: The plate was coated with purified Human IgA, IgG and IgM. Immunoglobulins were detected with serially diluted ARG21914 Goat anti-Human IgA antibody, F(ab')<sub>2</sub> fragment (FITC) (pre-adsorbed).