

## ARG21980 anti-CD4 antibody [GK1.5] (Biotin)

Package: 100 µg  
Store at: 4°C

### Summary

|                     |  |
|---------------------|--|
| Product Description | Biotin-conjugated Rat Monoclonal antibody [GK1.5] recognizes CD4                         |
| Tested Reactivity   | Ms   |
| Tested Application  | BL, Depletion, FACS, ICC/IF, IHC-Fr  |
| Specificity         | Mouse CD4  |
| Host                | Rat  |
| Clonality           | Monoclonal   |
| Clone               | GK1.5  |
| Isotype             | IgG2b, kappa   |
| Target Name         | CD4  |
| Species             | Mouse  |
| Immunogen           | Mouse CTL clone V4   |
| Conjugation         | Biotin   |
| Alternate Names     | CD4mut; CD antigen CD4; T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3 |

### Application Instructions

| Application table | Application | Dilution                     |
|-------------------|-------------|------------------------------|
|                   | BL          | Assay-dependent              |
|                   | Depletion   | Assay-dependent              |
|                   | FACS        | < 1 µg/10 <sup>6</sup> cells |
|                   | ICC/IF      | Assay-dependent              |
|                   | IHC-Fr      | Assay-dependent              |

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties


|                     |  |
|---------------------|--|
| Form                | Liquid   |
| Buffer              | PBS and 0.1% Sodium azide.   |
| Preservative        | 0.1% Sodium azide  |
| Concentration       | 0.5 mg/ml  |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

|                |  |
|----------------|--|
| Database links | <a href="#">GeneID: 12504 Mouse</a><br><a href="#">Swiss-port # P06332 Mouse</a>   |
| Gene Symbol    | CD4  |
| Gene Full Name | CD4 antigen  |
| Background     | CD4 is a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigens and is also a receptor for the human immunodeficiency virus. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene. [provided by RefSeq, Aug 2010]   |
| Function       | CD4 is an integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex. The antigens presented by class II peptides are derived from extracellular proteins while class I peptides are derived from cytosolic proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class II presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of T-helper cells. In other cells such as macrophages or NK cells, plays a role in differentiation/activation, cytokine expression and cell migration in a TCR/LCK-independent pathway. Participates in the development of T-helper cells in the thymus and triggers the differentiation of monocytes into functional mature macrophages. [UniProt] |
| Highlight      | Related products:<br><a href="#">CD4 antibodies</a> ; <a href="#">CD4 ELISA Kits</a> ; <a href="#">CD4 Duos / Panels</a> ; <a href="#">Anti-Rat IgG secondary antibodies</a> ;<br>Related news:<br><a href="#">New antibody panels and duos for Tumor immune microenvironment</a><br><a href="#">Tumor-Infiltrating Lymphocytes (TILs)</a>   |
| Research Area  | Developmental Biology antibody; Immune System antibody; Regulatory T cells Study antibody; T-cell infiltration Study antibody; Tumor-infiltrating Lymphocyte Study antibody  |
| Calculated Mw  | 51 kDa   |
| PTM            | Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.  |



ARG21980 anti-CD4 antibody [GK1.5] (Biotin) FACS image

Flow Cytometry: BALB/c Mouse splenocytes stained with ARG21980 anti-CD4 antibody [GK1.5] (Biotin) and ARG62941 anti-CD8a antibody [53-6.7] (PE).