

## ARG22679 anti-CD11b antibody [198]

Package: 50 µg  
Store at: -20°C

### Summary

|                     |  |
|---------------------|--|
| Product Description | Mouse Monoclonal antibody [198] recognizes CD11b<br>This antibody recognizes Rabbit CD11b cell surface glycoprotein, also known as the integrin alpha M chain and MAC-1. Mouse anti Rabbit CD11b antibody, clone 198 immunoprecipitates two proteins of molecular weight 165 kD and 95 kD from granulocytes. It recognizes monocytes, macrophages and neutrophils by flow cytometry and is thought to be against the homologue of Human CD11b. In immunohistochemistry good staining of macrophages is observed. |
| Tested Reactivity   | Rb   |
| Tested Application  | FACS, ICC/IF, IHC-Fr, IP   |
| Host                | Mouse  |
| Clonality           | Monoclonal   |
| Clone               | 198  |
| Isotype             | IgG1   |
| Target Name         | CD11b  |
| Species             | Rabbit   |
| Immunogen           | Rabbit adherent blood leucocytes.  |
| Conjugation         | Un-conjugated  |
| Alternate Names     | MAC1A; CR3A; CR-3 alpha chain; Cell surface glycoprotein MAC-1 subunit alpha; Integrin alpha-M; MAC-1; CD11 antigen-like family member B; Leukocyte adhesion receptor MO1; MO1A; SLEB6; Neutrophil adherence receptor; CD antigen CD11b; CD11B   |

### Application Instructions

| Application table | Application   | Dilution        |
|-------------------|---|-----------------|
|                   | FACS  | 1:100 - 1:200   |
|                   | ICC/IF  | Assay-dependent |
|                   | IHC-Fr  | Assay-dependent |
|                   | IP  | Assay-dependent |
| Application Note  | FACS: Use 10 µl of the suggested working dilution to label 10 <sup>6</sup> cells or 100 µl whole blood.<br>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. |                 |


### Properties

|              |                              |
|--------------|------------------------------|
| Form         | Liquid                       |
| Purification | Purification with Protein G. |
| Buffer       | PBS and 0.09% Sodium azide.  |

|                     |  |
|---------------------|--|
| Preservative        | 0.09% Sodium azide   |
| Concentration       | 1 mg/ml  |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note                | For laboratory research only, not for drug, diagnostic or other use.   |

## Bioinformation

|                |   |
|----------------|---|
| Gene Symbol    | ITGAM   |
| Gene Full Name | integrin, alpha M (complement component 3 receptor 3 subunit)   |
| Background     | CD11b (integrin alpha M chain): Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form a leukocyte-specific integrin referred to as macrophage receptor 1 ('Mac-1'), or inactivated-C3b (iC3b) receptor 3 ('CR3'). The alpha M beta 2 integrin is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009]  |
| Function       | CD11b: Integrin ITGAM/ITGB2 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PubMed:9558116, PubMed:20008295). It is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin ITGAM/ITGB2 is also a receptor for fibrinogen, factor X and ICAM1. It recognizes P1 and P2 peptides of fibrinogen gamma chain. Regulates neutrophil migration (PubMed:28807980). In association with beta subunit ITGB2/CD18, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed:21193407). May regulate phagocytosis-induced apoptosis in extravasated neutrophils. May play a role in mast cell development. Required with TYROBP/DAP12 in microglia to control production of microglial superoxide ions which promote the neuronal apoptosis that occurs during brain development. [UniProt] |
| Highlight      | Related products:<br><a href="#">CD11b antibodies: CD11b Duos / Panels: Anti-Mouse IgG secondary antibodies:</a><br>Related news:<br><a href="#">New antibody panels and duos for Tumor immune microenvironment</a><br><a href="#">Anti-SerpinB9 therapy, a new strategy for cancer therapy</a>   |
| Research Area  | MDSC Marker antibody; Myeloid-derived suppressor cell antibody  |
| Calculated Mw  | 127 kDa   |



ARG22679 anti-CD11b antibody [198] FACS image

Flow Cytometry: Rabbit peripheral blood granulocytes stained with ARG22679 anti-CD11b antibody [198].