

**ARG22858**  
anti-CD169 / Siglec 1 antibody [3B11/11] (FITC)Package: 50 µg  
Store at: 4°C

### Summary

Product Description	FITC-conjugated Mouse Monoclonal antibody [3B11/11] recognizes CD169 / Siglec 1
Tested Reactivity	Pig
Tested Application	FACS
Host	Mouse
Clonality	Monoclonal
Clone	3B11/11
Isotype	IgG1
Target Name	CD169 / Siglec 1
Species	Pig
Immunogen	Porcine alveolar macrophages.
Conjugation	FITC
Alternate Names	CD169; Siglec-1; dJ1009E24.1; Sialic acid-binding Ig-like lectin 1; SIGLEC-1; CD antigen CD169; SN; Sialoadhesin

### Application Instructions

Application table	Application	Dilution
	FACS	Neat

**Application Note** FACS: Use 10 µl of the suggested working dilution to 10<sup>6</sup> cells in 100 µl.  
\* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS, 0.09% Sodium azide and 1% BSA
Preservative	0.09% Sodium azide
Stabilizer	1% BSA
Concentration	0.1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Gene Symbol	SIGLEC1
Gene Full Name	sialic acid binding Ig-like lectin 1, sialoadhesin
Background	This gene encodes a member of the immunoglobulin superfamily. The encoded protein is a lectin-like adhesion molecule that binds glycoconjugate ligands on cell surfaces in a sialic acid-dependent manner. It is a type I transmembrane protein expressed only by a subpopulation of macrophages and is involved in mediating cell-cell interactions. Alternative splicing produces a transcript variant encoding an isoform that is soluble rather than membrane-bound; however, the full-length nature of this variant has not been determined. [provided by RefSeq, Jul 2008]
Function	Acts as an endocytic receptor mediating clathrin dependent endocytosis. Macrophage-restricted adhesion molecule that mediates sialic-acid dependent binding to lymphocytes, including granulocytes, monocytes, natural killer cells, B-cells and CD8 T-cells. Preferentially binds to alpha-2,3-linked sialic acid (By similarity). Binds to SPN/CD43 on T-cells (By similarity). May play a role in hemopoiesis. [UniProt]
Calculated Mw	183 kDa