

Summary

## ARG22966 anti-CD151 antibody [11G5a] (FITC)

Package: 50 μg Store at: 4°C

| Sammary             |  |
|---------------------|--|
| Product Description | FITC-conjugated Mouse Monoclonal antibody [11G5a] recognizes CD151<br>Mouse anti Human CD151 antibody, clone 11G5a recognizes the human CD151 cell surface antigen,<br>also known as PETA-3. CD151 is expressed by epithelial cells, endothelial cells, platelets,<br>megakaryocytes, monocytes and in the renal glomeruli and proximal and distal tubules. CD151 is not<br>expressed by lymphocytes or granulocytes. More recently CD151 has also been shown to be expressed<br>by erythrocytes, and to carry the MER2 blood group antigen (Crew et al. 2004). It should be noted that<br>CD151 is very closely associated with the alpha3 beta1 integrin in cells and co-immunoprecipitation<br>may occur even in quite stringent conditions (Yauch. et al. 1998). |
| Tested Reactivity   | Hu   |
| Tested Application  | FACS, ICC/IF, IHC-Fr   |
| Host                | Mouse  |
| Clonality           | Monoclonal   |
| Clone               | 11G5a  |
| lsotype             | lgG1   |
| Target Name         | CD151  |
| Species             | Human  |
| Conjugation         | FITC   |
| Alternate Names     | SFA1; Tetraspanin-24; GP27; Tspan-24; MER2; TSPAN24; PETA-3; RAPH; Membrane glycoprotein SFA-1;<br>CD151 antigen; CD antigen CD151; Platelet-endothelial tetraspan antigen 3   |

## **Application Instructions**

| Application table | Application  | Dilution        |
|-------------------|--|-----------------|
|                   | FACS   | Neat - 1:10     |
|                   | ICC/IF   | Assay-dependent |
|                   | IHC-Fr   | Assay-dependent |
| Application Note  | FACS: Use 10 $\mu$ l of the suggested working dilution to label 10^6 cells or 100 $\mu$ l whole blood.<br>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. |                 |

## Properties

| Form         | Liquid                             |
|--------------|------------------------------------|
| Purification | Purification with Protein A.       |
| Buffer       | PBS, 0.09% Sodium azide and 1% BSA |
| Preservative | 0.09% Sodium azide                 |

| Stabilizer          | 1% BSA   |
|---------------------|--|
| Concentration       | 0.1 mg/ml  |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note                | For laboratory research only, not for drug, diagnostic or other use.   |

## **Bioinformation**

| Gene Symbol    | CD151   |
|----------------|---|
| Gene Full Name | CD151 molecule (Raph blood group)   |
| Background     | The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins and other transmembrane 4 superfamily proteins. It is involved in cellular processes including cell adhesion and may regulate integrin trafficking and/or function. This protein enhances cell motility, invasion and metastasis of cancer cells. Multiple alternatively spliced transcript variants that encode the same protein have been described for this gene. [provided by RefSeq, Jul 2008] |
| Function       | Essential for the proper assembly of the glomerular and tubular basement membranes in kidney.<br>[UniProt]  |
| Calculated Mw  | 28 kDa  |