

ARG23048 anti-CD3 antibody [KT3] (FITC)

Package: 100 µg
Store at: 4°C

Summary

Product Description	FITC-conjugated Rat Monoclonal antibody [KT3] recognizes CD3 CD3 antibody, clone KT3 recognizes the mouse CD3 antigen expressed by mature T cells. Clone KT3 may be used to trigger proliferation and cytotoxicity of CD3 positive cells.
Tested Reactivity	Ms
Tested Application	FACS, IHC-Fr
Host	Rat
Clonality	Monoclonal
Clone	KT3
Isotype	IgG2a
Target Name	CD3
Species	Mouse
Immunogen	CBAT6 thymocytes.
Conjugation	FITC

Application Instructions

Application table	Application	Dilution
	FACS	1:50 - 1:100
	IHC-Fr	Assay-dependent
Application Note	FACS: Use 10 µl of the suggested working dilution to label 10 ⁶ cells in 100 µl. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Unpurified.
Buffer	PBS, 0.09% Sodium azide and 1% BSA
Preservative	0.09% Sodium azide
Stabilizer	1% BSA
Concentration	0.1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformatics

Gene Symbol	CD3D, CD3E, CD3G, CD247
Gene Full Name	CD3 Delta Subunit Of T-Cell Receptor Complex, CD3 Epsilon Subunit Of T-Cell Receptor Complex, CD3 Gamma Subunit Of T-Cell Receptor Complex, CD247 Molecule
Background	CD3 (cluster of differentiation 3) is a protein complex and T cell co-receptor that is involved in activating both the cytotoxic T cell (CD8+ naive T cells) and T helper cells (CD4+ naive T cells). It is composed of four distinct chains. In mammals, the complex contains a CD3 γ chain, a CD3 δ chain, and two CD3 ϵ chains. These chains associate with the T-cell receptor (TCR) and the CD3-zeta (ζ -chain) to generate an activation signal in T lymphocytes. The TCR, CD3-zeta, and the other CD3 molecules together constitute the TCR complex.
Highlight	Related products: CD3 antibodies ; CD3 ELISA Kits ; CD3 Duos / Panels ; CD3 recombinant proteins ; Anti-Rat IgG secondary antibodies ; Related news: New antibody panels and duos for Tumor immune microenvironment Tumor-Infiltrating Lymphocytes (TILs) Exploring Antiviral Immune Response
Calculated Mw	20 kDa

Images



ARG23048 anti-CD3 antibody [KT3] (FITC) FACS image

Flow Cytometry: Figure A. RPE conjugated Mouse anti Mouse CD28 and FITC conjugated Rat IgG2a isotype control. Figure B. RPE conjugated Mouse anti Mouse CD28 and ARG23048 anti-CD3 antibody [KT3] (FITC). All experiments performed on murine splenocytes.