

## ARG23298 anti-Macrophages + Monocytes antibody [KUL01]

Package: 100 µg  
Store at: -20°C

### Summary

|                     |  |
|---------------------|--|
| Product Description | Mouse Monoclonal antibody [KUL01] recognizes Macrophages + Monocytes<br>Mouse anti Chicken Monocytes/Macrophages, clone KUL01 recognises chicken monocytes and macrophages. Clone KUL01 will also detect interdigitating cells and activated microglia but does not recognise Bu1+ve B-cells or CD3 +ve T-lymphocytes. |
| Tested Reactivity   | Chk  |
| Tested Application  | FACS, IHC-Fr, IHC-P  |
| Host                | Mouse  |
| Clonality           | Monoclonal   |
| Clone               | KUL01  |
| Isotype             | IgG1   |
| Target Name         | Macrophages + Monocytes  |
| Species             | Chicken  |
| Conjugation         | Un-conjugated  |

### Application Instructions

| Application table | Application | Dilution        |
|-------------------|-------------|-----------------|
|                   | FACS        | Assay-dependent |
|                   | IHC-Fr      | Assay-dependent |
|                   | IHC-P       | Assay-dependent |

|                  |  |
|------------------|--|
| Application Note | IHC-P: This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase.<br>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. |
|------------------|--|

### Properties

|                     |  |
|---------------------|--|
| Form                | Liquid   |
| Purification        | Purified by ion exchange chromatography.   |
| Buffer              | BBS and 0.09% Sodium azide.  |
| Preservative        | 0.09% Sodium azide   |
| Concentration       | 0.5 mg/ml  |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |

**Note**

For laboratory research only, not for drug, diagnostic or other use.