

## ARG23325 anti-CD130 / gp130 antibody [B-S12] (low endotoxin)

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Azide free and low endotoxin Mouse Monoclonal antibody [B-S12] recognizes CD130 / gp130
Tested Reactivity	Hu
Tested Application	FACS, FuncSt, IP, WB
Specificity	This antibody recognizes the Gp130, common subunit for IL-6, IL-11, OSM, LIF, CNTF, CT-1 receptors, a 130-140 kDa protein.
Host	Mouse
Clonality	Monoclonal
Clone	B-S12
Isotype	IgG1
Target Name	CD130 / gp130
Species	Human
Immunogen	Natural soluble gp130
Conjugation	Un-conjugated
Alternate Names	CDw130; CD130; CDW130; Interleukin-6 signal transducer; CD antigen CD130; IL-6RB; Membrane glycoprotein 130; GP130; Oncostatin-M receptor subunit alpha; IL-6R subunit beta; Interleukin-6 receptor subunit beta; gp130; IL-6 receptor subunit beta; IL-6R-beta

### Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	FuncSt	Assay-dependent
	IP	Assay-dependent
	WB	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification Note	Sterile-filtered through 0.22 µm and treated to remove endotoxins.
Buffer	PBS
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Gene Symbol	IL6ST
Gene Full Name	interleukin 6 signal transducer
Background	The protein encoded by this gene is a signal transducer shared by many cytokines, including interleukin 6 (IL6), ciliary neurotrophic factor (CNTF), leukemia inhibitory factor (LIF), and oncostatin M (OSM). This protein functions as a part of the cytokine receptor complex. The activation of this protein is dependent upon the binding of cytokines to their receptors. vIL6, a protein related to IL6 and encoded by the Kaposi sarcoma-associated herpesvirus, can bypass the interleukin 6 receptor (IL6R) and directly activate this protein. Knockout studies in mice suggest that this gene plays a critical role in regulating myocyte apoptosis. Alternatively spliced transcript variants have been described. A related pseudogene has been identified on chromosome 17. [provided by RefSeq, May 2014]
Function	Signal-transducing molecule. The receptor systems for IL6, LIF, OSM, CNTF, IL11, CTF1 and BSF3 can utilize gp130 for initiating signal transmission. Binds to IL6/IL6R (alpha chain) complex, resulting in the formation of high-affinity IL6 binding sites, and transduces the signal. Does not bind IL6. May have a role in embryonic development (By similarity). The type I OSM receptor is capable of transducing OSM-specific signaling events. [UniProt]
Calculated Mw	104 kDa
PTM	Phosphorylation of Ser-782 down-regulates cell surface expression.  Heavily N-glycosylated (PubMed:11098061, PubMed:16335952, PubMed:19159218, PubMed:19139490, PubMed:11251120). Glycosylation is required for protein stability and localization in plasma membrane but not for ligand binding (PubMed:19915009). [UniProt]

## Images



ARG23325 anti-CD130 / gp130 antibody [B-S12] (low endotoxin)  
FACS image

Flow Cytometry: Eahy 926 cell line stained with ARG23325 anti-CD130 / gp130 antibody [B-S12] (low endotoxin).