

ARG23352 anti-CD9 antibody [TS9] (azide free)

Package: 100 µl
Store at: -20°C

Summary

Product Description	Azide free Mouse Monoclonal antibody [TS9] recognizes CD9
Tested Reactivity	Hu
Tested Application	FACS, IP, WB
Specificity	This antibody recognizes the p24 antigen, a 24 kDa protein.
Host	Mouse
Clonality	Monoclonal
Clone	TS9
Isotype	IgG1
Target Name	CD9
Species	Human
Immunogen	HeLa cell line
Conjugation	Un-conjugated
Alternate Names	Leukocyte antigen MIC3; BTCC-1; TSPAN-29; MIC3; Tetraspanin-29; p24; Cell growth-inhibiting gene 2 protein; CD9 antigen; MRP-1; DRAP-27; 5H9 antigen; CD antigen CD9; TSPAN29; Motility-related protein; Tspan-29

Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	IP	Assay-dependent
	WB	Assay-dependent
Application Note	WB: Under non-reducing conditions. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification Note	Sterile-filtered through 0.22 µm.
Buffer	PBS
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	CD9
Gene Full Name	CD9 molecule
Background	This gene encodes a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Tetraspanins are cell surface glycoproteins with four transmembrane domains that form multimeric complexes with other cell surface proteins. The encoded protein functions in many cellular processes including differentiation, adhesion, and signal transduction, and expression of this gene plays a critical role in the suppression of cancer cell motility and metastasis. [provided by RefSeq, Jan 2011]
Function	Involved in platelet activation and aggregation. Regulates paranodal junction formation. Involved in cell adhesion, cell motility and tumor metastasis. Required for sperm-egg fusion. [UniProt]
Highlight	Related products: CD9 antibodies ; CD9 ELISA Kits ; Anti-Mouse IgG secondary antibodies ; Related news: New antibodies for exosome isolation
Calculated Mw	25 kDa
PTM	Palmitoylated at a low, basal level in unstimulated platelets. The level of palmitoylation increases when platelets are activated by thrombin (in vitro). The protein exists in three forms with molecular masses between 22 and 27 kDa, and is known to carry covalently linked fatty acids. [UniProt]