

ARG23780 Goat anti-Mouse IgA + IgG + IgM antibody (Texas Red), pre-adsorbed

Package: 500 µg
Store at: 4°C

Summary

Product Description	Texas Red-conjugated Goat Polyclonal antibody recognizes Mouse IgA + IgG + IgM
Tested Reactivity	Ms
Tested Application	ELISA, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB
Specificity	Reacts with the heavy and light chains of mouse IgM, IgG1, IgG2a, IgG2b, IgG3 and IgA as demonstrated by ELISA, FLISA, and/or flow cytometry; minimal cross reactivity with human immunoglobulins
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgA + IgG + IgM
Species	Mouse
Conjugation	Texas Red

Application Instructions

Pre Adsorbed Human immunoglobulins and pooled sera.

Application table	Application	Dilution
	ELISA	Assay-dependent
	FACS	Assay-dependent
	FLISA	1:100 - 1:400
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
	IHC-P	Assay-dependent
	WB	Assay-dependent

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	1 mg/ml


Storage instruction

Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note


For laboratory research only, not for drug, diagnostic or other use.

Images



ARG23780 Goat anti-Mouse IgA + IgG + IgM antibody (Texas Red) (pre-adsorbed) ICC/IF image

Immunofluorescence: Human epithelial carcinoma cell line HEp-2 was stained with Mouse anti-Human CD44 (right) followed by ARG23780 Goat anti-Mouse IgA + IgG + IgM antibody (Texas Red) (pre-adsorbed) and DAPI.



ARG23780 Goat anti-Mouse IgA + IgG + IgM antibody (Texas Red) (pre-adsorbed) ICC/IF image

Immunofluorescence: U2-OS cells stimulated with tetradecanoylphorbol acetate and treated with MEK and p38 inhibitors were stained with anti-Bmi1 followed by ARG23780 Goat anti-Mouse IgA + IgG + IgM antibody (Texas Red) (pre-adsorbed).
