

Product datasheet

info@arigobio.com

ARG23878
Goat anti-Human IgG antibody, pre-adsorbed

Package: 500 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes Human IgG

Tested Reactivity Hu

Tested Application Depletion, ELISA, ELISPOT, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, IP, WB

Specificity The antibody reacts with the heavy chain of Human IgG. The antibody is pre-adsorbed with human IgM

and IgA so the antibody may not react to human IgM and IgA, but may react with IgG from other

species.

Host Goat

Clonality Polyclonal

Isotype IgG

Target Name IgG

Species Human

Immunogen Human IgG

Conjugation Un-conjugated

Application Instructions

Pre Adsorbed Human IgM and IgA.

Application table

Application	Dilution
Depletion	Assay-dependent
ELISA	Assay-dependent
ELISPOT	Assay-dependent
FACS	< 1 μg/10^6 cells
FLISA	Assay-dependent
ICC/IF	Assay-dependent
IHC-Fr	Assay-dependent
IHC-P	Assay-dependent
IP	Assay-dependent
WB	Assay-dependent
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	

Application Note

Properties

^{*} The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Form Liquid

Purification Affinity purification with immunogen.

Buffer BBS (pH 8.2)

Concentration 1 mg/ml


Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Images



ARG23878 Goat anti-Human IgG antibody (pre-adsorbed) ELISA image

ELISA: The plate was coated with purified Human IgG, IgM, and IgA. Immunoglobulins were detected with serially diluted ARG23878 Goat anti-Human IgG antibody (pre-adsorbed) followed by Mouse anti-Goat IgG (Fc) antibody (HRP).