

## ARG23880 Goat anti-Human IgA + IgG + IgM antibody (TRITC)

Package: 1 mg Store at: 4°C

## Summary

| Product Description | TRITC-conjugated Goat Polyclonal antibody recognizes Human IgA + IgG + IgM |  |
|---------------------|--|--|
| Tested Reactivity   | Hu   |  |
| Tested Application  | ELISA, FACS, FLISA, ICC/IF, WB   |  |
| Specificity         | Reacts with the heavy and light chains of Human IgG, IgM, and IgA.         |  |
| Host                | Goat   |  |
| Clonality           | Polyclonal   |  |
| lsotype             | IgG  |  |
| Target Name         | lgA + lgG + lgM  |  |
| Species             | Human  |  |
| Immunogen           | Human IgG, IgM, and IgA.   |  |
| Conjugation         | TRITC  |  |

## **Application Instructions**

| Application table | Application   | Dilution  |
|-------------------|---|---|
|                   | ELISA   | Assay-dependent   |
|                   | FACS  | Assay-dependent   |
|                   | FLISA   | 1:100 - 1:400   |
|                   | ICC/IF  | Assay-dependent   |
|                   | WB  | Assay-dependent   |
| Application Note  | * The dilutions indicate r<br>should be determined by | ecommended starting dilutions and the optimal dilutions or concentrations<br>the scientist. |

## **Properties**

| Form                | Liquid   |  |
|---------------------|--|--|
| Purification        | Affinity purification with immunogen.  |  |
| Buffer              | PBS and 0.1% Sodium azide.   |  |
| Preservative        | 0.1% Sodium azide  |  |
| Concentration       | 1 mg/ml  |  |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |  |