

ARG24073 Goat anti-Human IgG (Fc) antibody, pre-adsorbed

Package: 500 µg
Store at: -20°C

Summary

| | |
|---------------------|--|
| Product Description | Goat Polyclonal antibody recognizes Human IgG (Fc) |
| Tested Reactivity | Hu |
| Tested Application | ELISA, FACS |
| Specificity | The antibody reacts with the Fc region of human IgG. The antibody is pre-adsorbed with Human IgG Fab, IgM, IgA, serum proteins from rabbit, mouse, rat, bovine, horse, hamster, goat, sheep, chicken and guinea pig. The antibody may react with IgG from other species. |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | Human IgG (Fc) |
| Species | Human |
| Immunogen | Human IgG |
| Conjugation | Un-conjugated |

Application Instructions

Pre Adsorbed Human IgG Fab, IgM, IgA, serum proteins from rabbit, mouse, rat, bovine, horse, hamster, goat, sheep, chicken and guinea pig.

Application table

| Application | Dilution |
|-------------|-----------------|
| ELISA | 1 µg/ml |
| FACS | Assay-dependent |

Application Note

* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification | Affinity purification with immunogen. |
| Buffer | BBS (pH 8.2) |
| Concentration | 0.5 mg/ml |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |

Highlight

Related news:

[Comprehensive anti-Human secondary antibodies](#)