

ARG24079 Goat anti-Human IgG (Fc) antibody (Biotin), pre-adsorbed

Package: 500 µg
Store at: 4°C

Summary


Product Description	Biotin-conjugated Goat Polyclonal antibody recognizes Human IgG (Fc)
Tested Reactivity	Hu
Tested Application	ELISA, WB
Specificity	The antibody reacts with the Fc region of human IgG. The antibody is pre-adsorbed with Human IgG Fab, IgM and IgA, so the antibody may not react with Human IgG Fab, IgM and IgA, but may react with IgG from other species.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	Human IgG (Fc)
Species	Human
Immunogen	Human IgG
Conjugation	Biotin

Application Instructions

Pre Adsorbed	Human IgG Fab, IgM and IgA.	
Application table	Application	Dilution
	ELISA	1:5000 - 1:10000
	WB	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.



Dilution of Goat anti-Human IgG (Fc) antibody (Biotin) (pre-adsorbed)

ARG24079 Goat anti-Human IgG (Fc) antibody (Biotin) (pre-adsorbed) ELISA image

ELISA: The plate was coated with purified human IgG Fc, IgG Fab, IgM and IgA. Immunoglobulins were detected with serially diluted ARG24079 Goat anti-Human IgG (Fc) antibody (Biotin) (pre-adsorbed), followed by [ARG23912](#) Streptavidin (HRP).