

Product datasheet

info@arigobio.com

ARG24088 anti-CD4 antibody [CT7] (PE)

Package: 100 tests Store at: 4°C

Summary

Product Description PE-conjugated Mouse Monoclonal antibody [CT7] recognizes CD4

This antibody recognizes CD4 antigen present on T Helper/Inducer lymphocytes.

Tested Reactivity Gpig
Tested Application FACS

Host Mouse

Clonality Monoclonal

Clone CT7

Isotype IgG1

Target Name CD4

Species Guinea pig

Immunogen Guinea pig peritoneal T-cells.

Conjugation PE

Alternate Names CD4mut; CD antigen CD4; T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|---|
| | FACS | 10 μl to label 10^6 cells in 100 μl buffer or 100 μl whole Guinea Pig peripheral blood |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

Form Powder

Purification Purification with Protein G.

Buffer PBS, 0.09% Sodium azide, 1% BSA and 5% Sucrose.

Preservative 0.09% Sodium azide

Stabilizer 1% BSA and 5% Sucrose

Reconstitution Reconstitute with 1.0 ml of sterile distilled water

Storage instruction Store the lyophilized antibody at 2-8°C. After reconstitution, aliquot and store in the dark at 2-8°C. Keep

protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial

prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol CD4

Gene Full Name CD4 molecule

Background CD4 is a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex

class II antigenes and is also a receptor for the human immunodeficiency virus. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene. [provided by RefSeq, Aug 2010]

Function CD4 is an integral membrane glycoprotein that plays an essential role in the immune response and

serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex. The antigens presented by class II peptides are derived from extracellular proteins while class I peptides are derived from cytosolic proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class II presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of T-helper cells. In other cells such as macrophages or NK cells, plays a role in differentiation/activation, cytokine expression and cell migration in a TCR/LCK-independent pathway. Participates in the development of T-helper cells in the thymus and triggers the differentiation of monocytes into functional mature

macrophages. [UniProt]

Highlight Related products:

CD4 antibodies; CD4 ELISA Kits; CD4 Duos / Panels; Anti-Mouse IgG secondary antibodies;

Related news:

New antibody panels and duos for Tumor immune microenvironment

Tumor-Infiltrating Lymphocytes (TILs)


Research Area Developmental Biology antibody; Immune System antibody; Regulatory T cells Study antibody; T-cell

infiltration Study antibody; Tumor-infiltrating Lymphocyte Study antibody

Calculated Mw 51 kDa

PTM Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.

Images



ARG24088 anti-CD4 antibody [CT7] (PE) FACS image

Flow Cytometry: Stimulated Guinea Pig spleen cells stained with ARG24088 anti-CD4 antibody [CT7] (PE).