

ARG24100 Goat anti-Human Ig (Fab) antibody (FITC), pre-adsorbed

Package: 500 µg
Store at: 4°C

Summary


Product Description	FITC-conjugated Goat Polyclonal antibody recognizes Fab region of human IgG, IgM, and IgA
Tested Reactivity	Hu
Tested Application	ELISA, FLISA
Specificity	Reacts with the Fab region of Human Ig, IgM, and IgA. Pre-adsorbed with Human IgG Fc, so it does not react to Human IgG Fc region but may react with immunoglobulins from other species
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	Human IgG (Fab)
Species	Human
Immunogen	Human Ig
Conjugation	FITC

Application Instructions

Pre Adsorbed	Human IgG Fc.						
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>1:200 - 1:400</td></tr></tbody></table>	Application	Dilution	ELISA	Assay-dependent	FLISA	1:200 - 1:400
Application	Dilution						
ELISA	Assay-dependent						
FLISA	1:200 - 1:400						
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.						

Properties

Form	Liquid
Purification	Affinity chromatography on human pooled immunoglobulins covalently linked to agarose.
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.



ARG24100 Goat anti-Human Ig (Fab) antibody (FITC) (pre-adsorbed)
FLISA image

FLISA: The plate was coated with purified Human IgG Fab and IgG Fc. Immunoglobulins were detected with serially diluted ARG24100 Goat anti-Human Ig (Fab) antibody (FITC) (pre-adsorbed).