info@arigobio.com

# ARG24109
# Goat anti-Mouse IgG (H+L) antibody, F(ab')2 fragment, pre-adsorbed

Package: 250 µg
Store at: 4°C

## Summary

| | |
|---|---|
| Product Description | F(ab')2 fragment of Goat Polyclonal antibody recognizes Mouse IgG |
| Tested Reactivity | Ms |
| Tested Application | ELISA, ELISPOT, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB |
| Specificity | The antibody reacts with the heavy and light chains of mouse IgG1, IgG2a, IgG2b, IgG2c, and IgG3 and with the light chains of mouse IgM and IgA. |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | F(ab')2 IgG |
| Target Name | IgG |
| Species | Mouse |
| Immunogen | Mouse IgG |
| Conjugation | Un-conjugated |

## Application Instructions

| | |
|---|---|
| Pre Adsorbed | Mouse IgM and IgA, Human immunoglobulins and pooled sera |

Application table

| Application | Dilution |
|---|---|
| ELISA | Assay-dependent |
| ELISPOT | Assay-dependent |
| FACS | Assay-dependent |
| FLISA | Assay-dependent |
| ICC/IF | Assay-dependent |
| IHC-Fr | Assay-dependent |
| IHC-P | Assay-dependent |
| WB | Assay-dependent |

Application Note: * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

## Properties

| | |
|---|---|
| Form | Liquid |
| Buffer | PBS and 0.05% Sodium azide. |
| Preservative | 0.05% Sodium azide |

| | |
|---|---|
| Concentration | 0.5 mg/ml |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |

## Images




ARG24109 Goat anti-Mouse IgG (H+L) antibody, F(ab')2 fragment, pre-adsorbed ELISA image

ELISA: The plate was coated with purified Mouse IgM, IgG, and IgA and Human IgM, IgG, and IgA. Immunoglobulins were detected with serially diluted ARG24109 Goat anti-Mouse IgG (H+L) antibody, F(ab')2 fragment, pre-adsorbed.