

## ARG24115 anti-HYOU1 / ORP150 antibody [6E3-2C3]

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [6E3-2C3] recognizes HYOU1 / ORP150
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Mouse
Clonality	Monoclonal
Clone	6E3-2C3
Isotype	IgG2b
Target Name	HYOU1 / ORP150
Species	Human
Immunogen	Recombinant Full length HYOU1 / ORP150 Protein
Conjugation	Un-conjugated
Alternate Names	GRP-170; Hypoxia up-regulated protein 1; Grp170; ORP150; 150 kDa oxygen-regulated protein; 170 kDa glucose-regulated protein; HSP12A; ORP-150

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:100
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	140 kDa	


### Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS (pH 7.4), 50% Glycerol and 0.09% Sodium azide
Preservative	0.09% Sodium azide
Stabilizer	50% Glycerol
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

## Bioinformation


Gene Symbol	HYOU1
Gene Full Name	hypoxia up-regulated 1
Background	<p>The protein encoded by this gene belongs to the heat shock protein 70 family. This gene uses alternative transcription start sites. A cis-acting segment found in the 5' UTR is involved in stress-dependent induction, resulting in the accumulation of this protein in the endoplasmic reticulum (ER) under hypoxic conditions. The protein encoded by this gene is thought to play an important role in protein folding and secretion in the ER. Since suppression of the protein is associated with accelerated apoptosis, it is also suggested to have an important cytoprotective role in hypoxia-induced cellular perturbation. This protein has been shown to be up-regulated in tumors, especially in breast tumors, and thus it is associated with tumor invasiveness. This gene also has an alternative translation initiation site, resulting in a protein that lacks the N-terminal signal peptide. This signal peptide-lacking protein, which is only 3 amino acids shorter than the mature protein in the ER, is thought to have a housekeeping function in the cytosol. In rat, this protein localizes to both the ER by a carboxy-terminal peptide sequence and to mitochondria by an amino-terminal targeting signal. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2014]</p>
Function	<p>Has a pivotal role in cytoprotective cellular mechanisms triggered by oxygen deprivation. May play a role as a molecular chaperone and participate in protein folding. [UniProt]</p>
Calculated Mw	111 kDa
Cellular Localization	Endoplasmic reticulum lumen. [UniProt]

## Images



ARG24115 anti-HYOU1 / ORP150 antibody [6E3-2C3] ICC/IF image

Immunofluorescence: SK-N-BE stained with ARG24115 anti-HYOU1 / ORP150 antibody [6E3-2C3] at 1:100 dilution.



ARG24115 anti-HYOU1 / ORP150 antibody [6E3-2C3] WB image

Western blot: HEK293 stained with ARG24115 anti-HYOU1 / ORP150 antibody [6E3-2C3] at 1:1000 dilution.