

## ARG30342 Neuroendocrine Tumor IHC Antibody Panel

Package: 1 kit  
Store at: -20°C

### Component

Cat. No.	Component Name	Host clonality	Reactivity	Application	Package
ARG66347	anti-Ki-67 antibody [SQab1897]	Rabbit mAb	Hu	IHC-P	50 µl
ARG66772	anti-Chromogranin A antibody [SQab20205]	Rabbit mAb	Hu	IHC-P	50 µl
ARG66629	anti-Synaptophysin antibody [SQab19147]	Rabbit mAb	Hu, Ms, Rat	IHC-P, WB	50 µl
ARG66820	anti-CD56 / NCAM antibody [SQab20232]	Rabbit mAb	Hu	IHC-P	50 µl


### Summary

Target Name                      Neuroendocrine Tumor

### Properties


**Storage instruction**                      For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

**Note**    For laboratory research only, not for drug, diagnostic or other use.




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human tonsil tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human esophageal cancer tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human cervical cancer tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human diffuse large B-cell lymphoma stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human esophageal cancer tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).




ARG66347 anti-Ki-67 antibody [SQab1897] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human tonsil tissue stained with ARG66347 anti-Ki-67 antibody [SQab1897]. Antigen Retrieval: Heat mediation was performed using Tris/EDTA buffer (pH 9.0).



ARG66772 anti-Chromogranin A antibody [SQab20205] IHC-P image


Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human colon tissue. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). The tissue section was stained with ARG66772 anti-Chromogranin A antibody [SQab20205] at 18°C - 25°C for 30 minutes.



ARG66629 anti-Synaptophysin antibody [SQab19147] WB image


Western blot: 20 µg of Mouse brain and Rat brain lysates stained with ARG66629 anti-Synaptophysin antibody [SQab19147] at 1:500 dilution.





ARG66629 anti-Synaptophysin antibody [SQab19147] IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human glioma tissue stained with ARG66629 anti-Synaptophysin antibody [SQab19147]. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0).



ARG66820 anti-CD56 / NCAM antibody [SQab20232] IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human glioma tissue. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). The tissue section was stained with ARG66820 anti-CD56 / NCAM antibody [SQab20232].