

Product datasheet

info@arigobio.com

ARG40061 anti-NDUFB4 antibody

Package: 50 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes NDUFB4

Tested Reactivity Hu, Ms, Rat

Tested Application ICC/IF, IHC-P, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name NDUFB4

Species Human

Immunogen Recombinant fusion protein corresponding to aa. 1-90 of Human NDUFB4 (NP_001161803.1).

Conjugation Un-conjugated

Alternate Names NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 4; CI-B15; B15; NADH-ubiquinone

oxidoreductase B15 subunit; Complex I-B15

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse liver and HeLa	
Observed Size	15 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Gene Symbol NDUFB4

Gene Full Name NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 4, 15kDa

Background This gene encodes a non-catalytic subunit of the multisubunit NADH:ubiquinone oxidoreductase, the

first enzyme complex in the mitochondrial electron transport chain (complex I). Mammalian complex I is composed of 45 different subunits and transfers electrons from NADH to ubiquinone. [provided by

RefSeq, Dec 2009]

Function Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I),


that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be

ubiquinone. [UniProt]

Calculated Mw 15 kDa


Cellular Localization Mitochondrion inner membrane; Single-pass membrane protein; Matrix side. [UniProt]

Images



ARG40061 anti-NDUFB4 antibody WB image

Western blot: 25 μg of Mouse liver and HeLa cell lysates stained with ARG40061 anti-NDUFB4 antibody at 1:3000 dilution.



ARG40061 anti-NDUFB4 antibody WB image

Western blot: Extracts from normal (control) and NDUFB4 knockout (KO) HeLa cells, using ARG40061 anti-NDUFB4 antibody at 1:1000 dilution.