

ARG40457 anti-MXI1 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes MXI1
Tested Reactivity	Hu, Ms, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	MXI1
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 170-295 of Human MXI1 (NP_569157.2).
Conjugation	Un-conjugated
Alternate Names	Class C basic helix-loop-helix protein 11; Max interactor 1; bHLHc11; Max-interacting protein 1; MXI; MAD2; MXD2

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	26 kDa	

Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	MXI1
Gene Full Name	MAX interactor 1, dimerization protein
Background	Expression of the c-myc gene, which produces an oncogenic transcription factor, is tightly regulated in normal cells but is frequently deregulated in human cancers. The protein encoded by this gene is a transcriptional repressor thought to negatively regulate MYC function, and is therefore a potential tumor suppressor. This protein inhibits the transcriptional activity of MYC by competing for MAX, another basic helix-loop-helix protein that binds to MYC and is required for its function. Defects in this gene are frequently found in patients with prostate tumors. Three alternatively spliced transcripts encoding different isoforms have been described. Additional alternatively spliced transcripts may exist but the products of these transcripts have not been verified experimentally. [provided by RefSeq, Jul 2008]
Function	Transcriptional repressor. MXI1 binds with MAX to form a sequence-specific DNA-binding protein complex which recognizes the core sequence 5'-CAC[GA]TG-3'. MXI1 thus antagonizes MYC transcriptional activity by competing for MAX. [UniProt]
Calculated Mw	26 kDa
Cellular Localization	Nucleus. [UniProt]

Images


ARG40457 anti-MXI1 antibody WB image

Western blot: U 87MG, A549 and HepG2 stained with ARG40457 anti-MXI1 antibody at 1:3000 dilution.



ARG40457 anti-MXI1 antibody WB image

Western blot: Rat Brain stained with ARG40457 anti-MXI1 antibody at 1:3000 dilution.



ARG40457 anti-MXI1 antibody WB image

Western blot: Mouse Brain stained with ARG40457 anti-MXI1 antibody at 1:3000 dilution.

