

ARG40787 anti-REG3A antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes REG3A
Tested Reactivity	Hu
Tested Application	FACS, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	REG3A
Species	Human
Immunogen	KLH-conjugated synthetic peptide corresponding to aa. 129-163 of Human REG3A.
Conjugation	Un-conjugated
Alternate Names	HIP; PAP; HIP/PAP; Regenerating islet-derived protein 3-alpha; PAP-H; REG-III; Hepatointestinal pancreatic protein; PBCGF; Human proislet peptide; REG3; INGAP; Regenerating islet-derived protein III-alpha; REG-3-alpha; Reg III-alpha; PAP1; Pancreatitis-associated protein 1

Application Instructions

Application table	Application	Dilution
	FACS	1:25
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Purification with Protein A and immunogen peptide.
Buffer	PBS and 0.09% (W/V) Sodium azide.
Preservative	0.09% (W/V) Sodium azide.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	REG3A
Gene Full Name	regenerating islet-derived 3 alpha
Background	This gene encodes a pancreatic secretory protein that may be involved in cell proliferation or differentiation. It has similarity to the C-type lectin superfamily. The enhanced expression of this gene is observed during pancreatic inflammation and liver carcinogenesis. The mature protein also functions as an antimicrobial protein with antibacterial activity. Alternate splicing results in multiple transcript variants that encode the same protein.[provided by RefSeq, Nov 2014]
Function	Bactericidal C-type lectin which acts exclusively against Gram-positive bacteria and mediates bacterial killing by binding to surface-exposed carbohydrate moieties of peptidoglycan. Regulates keratinocyte proliferation and differentiation after skin injury via activation of EXTL3-PI3K-AKT signaling pathway. [UniProt]
Calculated Mw	19 kDa
PTM	Proteolytic processing by trypsin removes an inhibitory N-terminal propeptide and is essential for peptidoglycan binding and antibacterial activity. [UniProt]
Cellular Localization	Secreted. Note=Found in the apical region of pancreatic acinar cells. [UniProt]

Images



ARG40787 anti-REG3A antibody FACS image

Flow Cytometry: HepG2 cells stained with ARG40787 anti-REG3A antibody at 1:25 dilution (right histogram) or isotype control antibody (left histogram), followed by incubation with Alexa Fluor® 488 labelled secondary antibody.



ARG40787 anti-REG3A antibody WB image

Western blot: 35 µg of Human pancreas lysate stained with ARG40787 anti-REG3A antibody at 1:1000 dilution.