

ARG40949 anti-Desmoplakin I+II antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Desmoplakin I+II
Tested Reactivity	Hu, Ms, Rat
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Desmoplakin I+II
Species	Human
Immunogen	Synthetic peptide within aa. 2750-2850 of Human Desmoplakin (NP_004406.2).
Conjugation	Un-conjugated
Alternate Names	250/210 kDa paraneoplastic pemphigus antigen; Desmoplakin; DP; DP11; DCWHKTA; DPI

Application Instructions

Application table	Application	Dilution
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	300 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	DSP
Gene Full Name	desmoplakin
Background	Desmosomes are intercellular junctions that tightly link adjacent cells. Desmoplakin is an obligate component of functional desmosomes that anchors intermediate filaments to desmosomal plaques. The N-terminus of desmoplakin is required for localization to the desmosome and interacts with the N-terminal region of plakophilin 1 and plakoglobin. The C-terminus of desmoplakin binds with intermediate filaments. In the mid-region of desmoplakin, a coiled-coiled rod domain is responsible for homodimerization. Mutations in this gene are the cause of several cardiomyopathies and keratodermas as well as the autoimmune disease paraneoplastic pemphigus. [provided by RefSeq, Jul 2008]
Function	Major high molecular weight protein of desmosomes. Involved in the organization of the desmosomal cadherin-plakoglobin complexes into discrete plasma membrane domains and in the anchoring of intermediate filaments to the desmosomes. [UniProt]
Calculated Mw	332 kDa
PTM	Ser-2849 is probably phosphorylated by a cAMP-dependent protein kinase. Phosphorylation on Ser-2849 probably affects its association with epidermal, simple cytokeratins and VIM intermediate filaments. [UniProt]
Cellular Localization	Cell junction, desmosome. Cytoplasm, cytoskeleton. Cell membrane. Note=Innermost portion of the desmosomal plaque. Colocalizes with epidermal KRT5-KRT14 and simple KRT8-KRT18 keratins and VIM intermediate filaments network (PubMed:12802069). Localizes at the intercalated disk in cardiomyocytes (By similarity). [UniProt]

Images



ARG40949 anti-Desmoplakin I+II antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human gastric cancer tissue stained with ARG40949 anti-Desmoplakin I+II antibody at 1:100 dilution.