

## ARG41065 anti-Vitronectin antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes Vitronectin
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Vitronectin
Species	Human
Immunogen	Synthetic peptide derived from Human Vitronectin.
Conjugation	Un-conjugated
Alternate Names	Vitronectin; V75; VN; Serum-spreading factor; S-protein; VNT

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


## Bioinformation

---

Gene Symbol	VTN
Gene Full Name	vitronectin
Background	The protein encoded by this gene is a member of the pexin family. It is found in serum and tissues and promotes cell adhesion and spreading, inhibits the membrane-damaging effect of the terminal cytolytic complement pathway, and binds to several serpin serine protease inhibitors. It is a secreted protein and exists in either a single chain form or a clipped, two chain form held together by a disulfide bond. [provided by RefSeq, Jul 2008]
Function	<p>Vitronectin is a cell adhesion and spreading factor found in serum and tissues. Vitronectin interact with glycosaminoglycans and proteoglycans. Is recognized by certain members of the integrin family and serves as a cell-to-substrate adhesion molecule. Inhibitor of the membrane-damaging effect of the terminal cytolytic complement pathway.</p> <p>Somatomedin-B is a growth hormone-dependent serum factor with protease-inhibiting activity. [UniProt]</p>
Calculated Mw	54 kDa
PTM	<p>Sulfated on 2 tyrosine residues.</p> <p>N- and O-glycosylated.</p> <p>Phosphorylation on Thr-69 and Thr-76 favors cell adhesion and spreading.</p> <p>It has been suggested that the active SMB domain may be permitted considerable disulfide bond heterogeneity or variability, thus two alternate disulfide patterns based on 3D structures are described with 1 disulfide bond conserved in both.</p> <p>Phosphorylation sites are present in the extracellular medium. [UniProt]</p>
Cellular Localization	Secreted, extracellular space. [UniProt]

## Images

---



Human serum membrane

ARG41065 anti-Vitronectin antibody WB image

Western blot: Human serum membrane lysate stained with ARG41065 anti-Vitronectin antibody.