

## ARG41139 anti-TGFBI antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes TGFBI
Tested Reactivity	Hu, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	TGFBI
Species	Human
Immunogen	Synthetic peptide derived from Human TGFBI.
Conjugation	Un-conjugated
Alternate Names	CDGG1; LCD1; RGD-CAP; CSD2; CSD; Beta ig-h3; CSD1; Transforming growth factor-beta-induced protein ig-h3; RGD-containing collagen-associated protein; BIGH3; CDG2; CSD3; Kerato-epithelin; CDB1; EBMD

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Human fetal kidney	

### Properties


Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	TGFBI
Gene Full Name	transforming growth factor, beta-induced, 68kDa
Background	This gene encodes an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. This protein plays a role in cell-collagen interactions and may be involved in endochondrial bone formation in cartilage. The protein is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in this gene are associated with multiple types of corneal dystrophy. [provided by RefSeq, Jul 2008]
Function	Binds to type I, II, and IV collagens. This adhesion protein may play an important role in cell-collagen interactions. In cartilage, may be involved in endochondral bone formation. [UniProt]
Calculated Mw	75 kDa
PTM	Gamma-carboxylation is controversial. Gamma-carboxyglutamated; gamma-carboxyglutamate residues are formed by vitamin K dependent carboxylation; these residues may be required for binding to calcium (PubMed:18450759). According to a more recent report, does not contain vitamin K-dependent gamma-carboxyglutamate residues (PubMed:26273833).  The EMI domain contains 2 expected intradomain disulfide bridges (Cys-49-Cys85 and Cys-84-Cys-97) and one unusual interdomain disulfide bridge to the second FAS1 domain (Cys-74-Cys-339). This arrangement violates the predicted disulfide bridge pattern of an EMI domain. [UniProt]
Cellular Localization	Secreted. Secreted, extracellular space, extracellular matrix. Note=May be associated both with microfibrils and with the cell surface (PubMed:8077289). [UniProt]

## Images



ARG41139 anti-TGFBI antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human kidney stained with ARG41139 anti-TGFBI antibody.



Human fetal kidney

ARG41139 anti-TGFBI antibody WB image

Western blot: Human fetal kidney lysate stained with ARG41139 anti-TGFBI antibody.