

Product datasheet

info@arigobio.com

ARG41179 anti-ACTR1A + ACTR1B antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes ACTR1A + ACTR1B

Tested Reactivity Hu, Ms, Rat

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name ACTR1A + ACTR1B

Species Human

Immunogen Recombinant fusion protein corresponding to aa. 187-376 of Human ACTR1A (NP_005727.1).

Conjugation Un-conjugated

Alternate Names Alpha-centractin; Actin-RPV; Arp1A; Centractin; Centrosome-associated actin homolog; CTRN1; ARP1

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse spleen	
Observed Size	45 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol ACTR1A

Gene Full Name ARP1 actin-related protein 1 homolog A, centractin alpha (yeast)

Background This gene encodes a 42.6 kD subunit of dynactin, a macromolecular complex consisting of 10-11

subunits ranging in size from 22 to 150 kD. Dynactin binds to both microtubules and cytoplasmic dynein. It is involved in a diverse array of cellular functions, including ER-to-Golgi transport, the centripetal movement of lysosomes and endosomes, spindle formation, chromosome movement, nuclear positioning, and axonogenesis. This subunit is present in 8-13 copies per dynactin molecule, and

is the most abundant molecule in the dynactin complex. It is an actin-related protein, and is approximately 60% identical at the amino acid level to conventional actin. [provided by RefSeq, Jul

2008]

Function Component of a multi-subunit complex involved in microtubule based vesicle motility. It is associated


with the centrosome. [UniProt]

Calculated Mw 43 kDa


Cellular Localization Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.

Cytoplasm, cell cortex. [UniProt]

Images



Western blot: $25~\mu g$ of Mouse spleen lysate stained with ARG41179 anti-ACTR1A + ACTR1B antibody at 1:1000 dilution.



Mouse spleen