

ARG41483 anti-Glycophorin C antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Glycophorin C
Tested Reactivity	Hu, Ms
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Glycophorin C
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 1-128 of Human Glycophorin C (NP_002092.1).
Conjugation	Un-conjugated
Alternate Names	Glycophorin-D; CD236R; GPD; Glycophorin-C; Glycoprotein beta; Glycoconnectin; GPC; PAS-2; CD antigen CD236; GE; CD236; Sialoglycoprotein D; GYPD; PAS-2'

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HepG2	
Observed Size	~ 34 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	GYPC
Gene Full Name	glycophorin C (Gerbich blood group)
Background	Glycophorin C (GYPC) is an integral membrane glycoprotein. It is a minor species carried by human erythrocytes, but plays an important role in regulating the mechanical stability of red cells. A number of glycophorin C mutations have been described. The Gerbich and Yus phenotypes are due to deletion of exon 3 and 2, respectively. The Webb and Duch antigens, also known as glycophorin D, result from single point mutations of the glycophorin C gene. The glycophorin C protein has very little homology with glycophorins A and B. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Feb 2012]
Function	This protein is a minor sialoglycoprotein in human erythrocyte membranes. The blood group Gerbich antigens and receptors for Plasmodium falciparum merozoites are most likely located within the extracellular domain. Glycophorin-C plays an important role in regulating the stability of red cells. [UniProt]
Calculated Mw	14 kDa
PTM	O-glycosylated with core 1 or possibly core 8 glycans. [UniProt]
Cellular Localization	Cell membrane; Single-pass type III membrane protein. Note=Linked to the membrane via band 4.1. [UniProt]

Images



ARG41483 anti-Glycophorin C antibody WB image

Western blot: 25 µg of HepG2 cell lysate stained with ARG41483 anti-Glycophorin C antibody at 1:1000 dilution.