

Product datasheet

info@arigobio.com

ARG41691 anti-RBMS1 antibody

Package: 50 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes RBMS1

Tested Reactivity Hu

Predict Reactivity Hu, Ms, Rat, Cow, Dog, Goat, Gpig, Hrs, Rb

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name RBMS1
Species Human

Immunogen Synthetic peptide around the C-terminal region of Human RBMS1. (within the following region:

TYMPATSAMQGAYLPQYAHMQTTAVPVEEASGQQQVAVETSNDHSPYTFQ)

Conjugation Un-conjugated

Alternate Names HCC-4; SCR2; YC1; RNA-binding motif, single-stranded-interacting protein 1; C2orf12; Single-stranded

DNA-binding protein MSSP-1; MSSP-1; MSSP-2; MSSP-3; Suppressor of CDC2 with RNA-binding

motif 2

Application Instructions

Predict Reactivity Note Predicted Homology Based On Immunogen Sequence: Cow: 93%; Dog: 100%; Goat: 93%; Guinea pig:

100%; Horse: 100%; Mouse: 100%; Rabbit: 100%; Rat: 100%

Application table Application Dilution

WB 2 - 4 μg/ml

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations

should be determined by the scientist.

Positive Control Jurkat

Observed Size ~ 47 kDa

Properties

Form Liquid

Purification Purification with Protein A.

Buffer PBS, 0.09% (w/v) Sodium azide and 2% Sucrose.

Preservative 0.09% (w/v) Sodium azide

Stabilizer 2% Sucrose

Concentration Batch dependent: 0.5 - 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol RBMS1

Gene Full Name RNA binding motif, single stranded interacting protein 1

Background This gene encodes a member of a small family of proteins which bind single stranded DNA/RNA. These

proteins are characterized by the presence of two sets of ribonucleoprotein consensus sequence (RNP-CS) that contain conserved motifs, RNP1 and RNP2, originally described in RNA binding proteins, and required for DNA binding. These proteins have been implicated in such diverse functions as DNA replication, gene transcription, cell cycle progression and apoptosis. Several transcript variants, resulting from alternative splicing and encoding different isoforms, have been described. A pseudogene

for this locus is found on chromosome 12. [provided by RefSeq, Feb 2009]

Function Single-stranded DNA binding protein that interacts with the region upstream of the MYC gene. Binds


specifically to the DNA sequence motif 5'-[AT]CT[AT][AT]T-3'. Probably has a role in DNA replication.

[UniProt]

Calculated Mw 45 kDa

Cellular Localization Nucleus. [UniProt]

Images



ARG41691 anti-RBMS1 antibody WB image

Western blot: Jurkat cell lysate stained with ARG41691 anti-RBMS1 antibody at 2.5 $\mu g/ml$ dilution.