

ARG41989 anti-MEK3 + MEK6 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes MEK3 + MEK6
Tested Reactivity	Hu, Ms, Rat
Tested Application	FACS, ICC/IF, IHC-P, IP, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	MEK3 + MEK6
Species	Human
Immunogen	Synthetic peptide of Human MEK3 / MEK6.
Conjugation	Un-conjugated
Alternate Names	<p>MEK3: SAPK kinase 2; MEK 3; MAPKK 3; Stress-activated protein kinase kinase 2; PRKMK3; EC 2.7.12.2; MAPK/ERK kinase 3; MAPKK3; SAPKK2; SAPKK-2; MAP kinase kinase 3; MKK3; Dual specificity mitogen-activated protein kinase kinase 3</p> <p>MEK6: SAPK kinase 3; MEK 6; MAPKK 6; MAPK/ERK kinase 6; EC 2.7.12.2; PRKMK6; SAPKK3; MAPKK6; SAPKK-3; Stress-activated protein kinase kinase 3; MKK6; MAP kinase kinase 6; Dual specificity mitogen-activated protein kinase kinase 6</p>

Application Instructions

Application table	Application	Dilution
	FACS	1:50
	ICC/IF	1:100 - 1:500
	IHC-P	1:50 - 1:200
	IP	1:50
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	
Observed Size	37 and 39 kDa	

Properties

Form	Liquid
Purification	Affinity purified.

Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	MAP2K3; MAP2K6
Gene Full Name	mitogen-activated protein kinase kinase 3 mitogen-activated protein kinase kinase 6
Background	<p>MEK3: The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is activated by mitogenic and environmental stress, and participates in the MAP kinase-mediated signaling cascade. It phosphorylates and thus activates MAPK14/p38-MAPK. This kinase can be activated by insulin, and is necessary for the expression of glucose transporter. Expression of RAS oncogene is found to result in the accumulation of the active form of this kinase, which thus leads to the constitutive activation of MAPK14, and confers oncogenic transformation of primary cells. The inhibition of this kinase is involved in the pathogenesis of Yersinia pseudotuberculosis. Multiple alternatively spliced transcript variants that encode distinct isoforms have been reported for this gene. [provided by RefSeq, Jul 2008]</p> <p>MEK6: This gene encodes a member of the dual specificity protein kinase family, which functions as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. As an essential component of p38 MAP kinase mediated signal transduction pathway, this gene is involved in many cellular processes such as stress induced cell cycle arrest, transcription activation and apoptosis. [provided by RefSeq, Jul 2008]</p>
Function	Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. With MAP3K3/MKK3, catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in the MAP kinases p38 MAPK11, MAPK12, MAPK13 and MAPK14 and plays an important role in the regulation of cellular responses to cytokines and all kinds of stresses. Especially, MAP2K3/MKK3 and MAP2K6/MKK6 are both essential for the activation of MAPK11 and MAPK13 induced by environmental stress, whereas MAP2K6/MKK6 is the major MAPK11 activator in response to TNF. MAP2K6/MKK6 also phosphorylates and activates PAK6. The p38 MAP kinase signal transduction pathway leads to direct activation of transcription factors. Nuclear targets of p38 MAP kinase include the transcription factors ATF2 and ELK1. Within the p38 MAPK signal transduction pathway, MAP3K6/MKK6 mediates phosphorylation of STAT4 through MAPK14 activation, and is therefore required for STAT4 activation and STAT4-regulated gene expression in response to IL-12 stimulation. The pathway is also crucial for IL-6-induced SOCS3 expression and down-regulation of IL-6-mediated gene induction; and for IFNG-dependent gene transcription. Has a role in osteoclast differentiation through NF-kappa-B transactivation by TNFSF11, and in endochondral ossification and since SOX9 is another likely downstream target of the p38 MAPK pathway. MAP2K6/MKK6 mediates apoptotic cell death in thymocytes. Acts also as a regulator for melanocytes dendricity, through the modulation of Rho family GTPases. [UniProt]
Calculated Mw	MEK3: 39 kDa MEK6: 37 kDa
PTM	<p>MEK3: Autophosphorylated. Phosphorylation on Ser-218 and Thr-222 by MAP kinase kinase kinases regulates positively the kinase activity (PubMed:8622669). Phosphorylated by TAOK2 (PubMed:11279118).</p> <p>Yersinia yopJ may acetylate Ser/Thr residues, preventing phosphorylation and activation, thus blocking the MAPK signaling pathway. [UniProt]</p> <p>MEK6: Weakly autophosphorylated. Phosphorylated at Ser-207 and Thr-211 by the majority of M3Ks, such as MAP3K5/ASK1, MAP3K1/MEKK1, MAP3K2/MEKK2, MAP3K3/MEKK3, MAP3K4/MEKK4, MAP3K7/TAK1, MAP3K11/MLK3 and MAP3K17/TAOK2.</p>

Acetylation of Ser-207 and Thr-211 by Yersinia yopJ prevents phosphorylation and activation, thus blocking the MAPK signaling pathway. [UniProt]

Cellular Localization


MEK6: Nucleus. Cytoplasm. Cytoplasm, cytoskeleton. Note=Binds to microtubules. [UniProt]

Images




ARG41989 anti-MEK3 + MEK6 antibody ICC/IF image

Immunofluorescence: HeLa cells stained with ARG41989 anti-MEK3 + MEK6 antibody.



ARG41989 anti-MEK3 + MEK6 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse spleen tissue stained with ARG41989 anti-MEK3 + MEK6 antibody.



ARG41989 anti-MEK3 + MEK6 antibody WB image

Western blot: HeLa cell lysate stained with ARG41989 anti-MEK3 + MEK6 antibody.